

Corporate Medical Policy

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “**Notification**”

File Name: biomarkers_for_myocardial_infarction_and_chronic_heart_failure
Origination: 01/2019
Last Review: 07/2023

Effective Date 10/24/23

Description of Procedure or Service

Cardiac biomarkers are the biochemical markers released in blood from injured myocardial tissue in both acute and chronic conditions, such as myocardial infarction or heart failure. They become elevated in blood after a certain period and can be measured. Examples of cardiac biomarkers commonly used in the acute clinical setting include Troponin and Creatine Kinase MB isoenzyme (CKMB) (Thygesen et al., 2007). Others, such as Suppression of tumorigenicity 2 (ST2), can serve in long-term as markers of cardiomyocyte stress and fibrosis for risk stratification of patients with a wide spectrum of cardiovascular diseases (Bayes-Genis et al., 2015).

Related Policies

Cardiovascular Disease Risk Assessment AHS – G2050

******Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.***

Policy

BCBSNC will provide coverage for biomarkers for myocardial infarction and chronic heart failure when it is determined the medical criteria or reimbursement guidelines below are met.

Benefits Application

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore member benefit language should be reviewed before applying the terms of this medical policy.

When Biomarkers for Myocardial Infarction and Chronic Heart Failure is covered

For individuals presenting with signs and symptoms of acute coronary syndrome (see Note 1), reimbursement is allowed for **quantitative** measurement of cardiac troponin (troponin T or I) for the diagnosis of myocardial infarction (MI) (when tested at an outpatient facility capable of performing an adequate clinical MI evaluation) up to four times within the first 72 hours following initial presentation.

When Biomarkers for Myocardial Infarction and Chronic Heart Failure is not covered

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

1. For individuals presenting with signs and symptoms of acute coronary syndrome (see Note 1), reimbursement is not allowed for measurement of the following cardiac biomarkers for the diagnosis and/or prognosis of MI:
 - a. Aspartate aminotransferase (AST/SGOT).
 - b. Cardiac creatine kinase isoenzyme MB (CKMB).
 - c. Creatine kinase (CK).
 - d. Creatine kinase isoenzymes.
 - e. Lactate dehydrogenase (LD, LDH).
 - f. Myoglobin.
2. For individuals presenting with signs and symptoms of acute coronary syndrome (see Note 1), reimbursement is not allowed for measurement of cardiac biomarkers in an outpatient setting which is not capable of performing adequate clinical MI evaluation (e.g., independent lab or physician’s office).
3. For all situations, reimbursement is not allowed for **qualitative** measurement of cardiac troponin (troponin T or I).
4. For individuals presenting with signs and symptoms of acute coronary syndrome (see Note 1), reimbursement is not allowed for measurement of the following cardiac biomarkers for the diagnosis and/or prognosis of MI:
 - a. Copeptin.
 - b. Troponin C.
 - c. C-reactive protein.
 - d. Heart-type fatty acid binding protein (H-FABP).
 - e. Any other cardiac biomarkers not listed above.
5. For all situations in the outpatient setting, reimbursement is not allowed for analysis of ST2 and/or its isoforms (e.g., Presage ST2).

NOTES:

Note 1: *Acute Coronary Syndrome/Myocardial Infarction Common Signs and Symptoms (Reeder, 2022):*

- Ischemic chest pain with radiation to an upper extremity, radiation to both arms, and pain associated with diaphoresis or with nausea and vomiting.
- Squeezing, tightness, pressure, constriction, crushing, strangling, burning, heartburn, fullness in the chest, band-like sensation, knot in the center of the chest, lump in throat, ache, heavy weight on chest and toothache (when there is radiation to the lower jaw).
- Ischemic pain often radiates to other parts of the body including the upper abdomen (epigastrium), shoulders, arms (upper and forearm), wrist, fingers, neck and throat, lower jaw and teeth (but not upper jaw), and not infrequently to the back (specifically the interscapular region).
- Shortness of breath, belching, nausea, indigestion, vomiting, diaphoresis, dizziness, lightheadedness, clamminess, and fatigue.

Atypical Signs and Symptoms (Reeder, 2022):

Dyspnea alone, weakness, nausea and/or vomiting, epigastric pain or discomfort, palpitations, syncope, or cardiac arrest.

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

Policy Guidelines

Background

Acute coronary syndromes (ACS) represent continuous events starting with angina, reversible injury, progressing to unstable angina, are frequently associated with minor myocardial damage, and myocardial infarction (MI) that results in extensive tissue necrosis (Thygesen et al., 2007). Patients with ACS usually present with chest pain and associated signs and symptoms. These patients are subdivided into two major categories based on the 12-lead electrocardiogram (ECG). If an ST-segment elevation is observed on the ECG, it is indicative of acute ST-elevation myocardial infarction (STEMI) type of ACS. If the ECG shows ST-segment depression, T-wave changes, or no ECG abnormalities, it is indicative of non-ST elevation myocardial infarction (NSTEMI) and unstable angina. ACS is complex. However, the most common cause is atherosclerotic coronary artery disease with rupture of atherosclerotic plaque (Amsterdam et al., 2014). The first documented definition of acute MI was established in 1979 by the World Health Organization (WHO). It included in the criteria for MI diagnosis the recommendation to use the rise or fall patterns of cardiac biomarkers, such as creatine kinase (CK), creatine kinase’s MB isoenzyme (CK-MB), lactate dehydrogenase (LDH) or aspartate aminotransferase (AST) activities (WHO, 1979). Since then, other societies have proposed their own criteria for diagnosis. The third universal definition of MI includes typical clinical symptoms, suggestive ECG changes, or imaging evidence of new loss of viable myocardium or new regional wall abnormality with a rise and/or fall of cardiac biomarkers (Thygesen et al., 2012). Nonetheless, the universal criteria are being refined by cardiovascular societies and will likely change with scientific progress and better understanding of MI pathophysiology.

Myocardial infarction results in cardiac injury and extensive tissue necrosis. The cellular membranes become compromised and release structural proteins and other macromolecules into cardiac interstitial, called cardiac biomarkers. The levels of these cardiac biomarkers in blood will rise and fall with time after MI (Thygesen et al., 2007). The first cardiac biomarker, aspartate aminotransferase (AST), was used for MI diagnosis in 1954. AST is present in human tissues as two isoenzymes: cytoplasmic and mitochondrial. AST is a non-specific biomarker, and its activity could also be elevated in other conditions, such as hepatic congestion secondary to congestive heart failure. Since then, other cardiac biomarkers were used as an aid in diagnosis of MI, but due to their non-specificity and other reasons, many of them are no longer used in clinical practice or their use remains very limited (Danese & Montagnana, 2016). The most common cardiac biomarkers and their characteristics are summarized in the table from (Danese & Montagnana, 2016):

Table 1 Characteristics of AMI biomarkers

Biomarker	First assay development, year	Molecular weight, Da	Kinetics			Sensitivity for myocyte necrosis	Specificity for myocyte necrosis
			First detection, hours	Maximum value, hours	Return to normal values, days		
AST	1954	105,000	3–4	15–28	5	++	+
LDH	1955	140,000	5–10	60–144	12	++	+
CK total enzyme activity	1960	83,000	3–9	10–20	3	++	+
CK-MB activity	1972	83,000	3–8	10–20	3	++	++
Myoglobin	1978	17,800	1–3	4–7	1–1.5	+++	+
CK-MB mass	1985	83,000	3–12	12–18	2–3	+++	+++
cTnI	1987	23,900	3–7	10–20	10	++++	++++
cTnT	1989	37,000	3–8	15–120	14	++++	++++

AST, aspartate aminotransferase; LDH, lactate dehydrogenase; CK, creatine kinase; CK-MB, creatine kinase MB isoenzyme; cTnI, cardiac troponin I; cTnT, cardiac troponin T.

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

Heart failure (HF) is a complex clinical syndrome resulting from any structural or functional impairment of ventricular filling or ejection of blood, including disorders of the pericardium, myocardium, endocardium, heart valves, great vessels, or certain metabolic abnormalities (Colucci, 2022). Most patients with HF have symptoms due to impaired left ventricular (LV) myocardial function (Colucci & Dunlay, 2022; Yancy et al., 2013). The most common symptoms of HF are dyspnea and fatigue, which may limit exercise tolerance and fluid retention. Some patients have exercise intolerance but little evidence of fluid retention, whereas others complain primarily of edema, dyspnea, or fatigue (Colucci & Dunlay, 2022). Heart failure is often a progressive condition, beginning with predisposing factors and leading to the development and worsening of clinical illness (Colucci, 2021; Colucci & Dunlay, 2022).

Lactate Dehydrogenase (LDH, also known as LD)

Lactate dehydrogenase is a cytoplasmic enzyme present in many different tissues, such as skeletal muscle, liver, heart, kidney, and red blood cells. Five isoenzymes have been identified by gel electrophoresis and other techniques (Marshall et al., 1991). The heart isoenzymes, LD1 and LD2, have activity increases in blood five to ten hours after MI symptoms onset and remains elevated for up to ten days (Danese & Montagnana, 2016). LD has poor specificity for cardiac tissue and is generally not recommended as a biomarker for the diagnosis of MI (Amsterdam et al., 2014; Jaffe & Morrow, 2021).

Myoglobin

Myoglobin is an oxygen-binding, cytoplasmic, heme protein. It is one of the first cardiac biomarkers measurable in the serum that appears between one and three hours after MI symptoms onset. Myoglobin is present in skeletal and cardiac muscles and is cleared by the kidneys (Vaidya, 1994). Its clinical utility is limited by its poor specificity. The main reason of using myoglobin in a clinical setting was its sensitivity for MI (Danese & Montagnana, 2016); but with appearance of sensitive troponin assays, myoglobin use offers little advantage for the diagnosis of MI (Eggers et al., 2004; Kavsak et al., 2007). Currently, there are no recommendations for myoglobin to be used in the diagnosis of MI (Amsterdam et al., 2014), and its use as cardiac biomarker is discouraged (Amsterdam et al., 2014; Jaffe & Morrow, 2021).

Creatine Kinase (CK) Isoenzymes and Isoform MB (CKMB)

The cytosolic enzyme creatine kinase (CK), formerly known as creatine phosphokinase (Danese & Montagnana, 2016), is present as three cytosolic isoenzymes and one mitochondrial isoenzyme. These isoenzymes are dimers of M (muscle) and B (brain) chains that exist in three combinations: MM, MB and BB (Bessman & Carpenter, 1985). The CKMM is predominant in both heart and skeletal muscle, but CKMB is more specific for the myocardium. The total CK activity could be detected in blood 3-9 hours after MI, but it reaches the maximum levels in blood in 10-20 hours and returns to normal in about 72 hours (Penttila et al., 2000). The measurement of total CK activity is not specific to MI because it also increases in liver, biliary tract, kidneys, and skeletal muscle disease, and its measurement is problematic in older individuals with lower muscle mass (Dillon et al., 1982; Heller et al., 1983; Yusuf et al., 1987). CKMB mass (CKMB protein concentration measurements) was once the cardiac biomarker of the choice that replaced CK, CKMB activity, AST, and LDH (Danese & Montagnana, 2016). However, with arrival of cardiac troponin assays, the use of CKMB became less popular. Some clinicians advocate for the use of CKMB for diagnosis and prognosis of MI, but cardiac troponins have shown either equally reliable or superior results compared to CKMB; consequently, troponin is the recommended test for MI diagnosis now (Amsterdam et al., 2014; Jaffe & Morrow, 2021).

Engel and Rockson (2020) studied the use of CK-MB in early diagnosis of myocardial infarction within the first nine hours of the hospital stay. 528 patient charts with complaints of chest pain within the past year were studied. An enzymatic diagnosis was assigned if CK-MB exceeded the normal values. The diagnosis of each patient before 9 hours (early diagnosis) was compared to the ultimate diagnosis at 14-24 hours (final diagnosis). Of the 528 patients, 195 (36.9%) had an early MI diagnosis within 9 hours and 190 patients (97.4%) of these did have an ultimate diagnosis of MI. Therefore, the authors conclude

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS

– G2150 “Notification”

that “standard CK-MB measurements within 9 hours of arrival provided an accurate clinical assessment in > 99% of the cases” (Engel & Rockson, 2020).

Troponins

The regulatory protein troponin in the troponin complex is composed of three isoforms. Troponin C (TnC) is responsible for calcium binding and has no role to play as a cardiac biomarker. Troponin I (TnI) and Troponin T (TnT) are responsible for inhibition of ATPase activity and tropomyosin binding, respectively (Greaser & Gergely, 1971). Contrary to all previously used cardiac biomarkers, cardiac troponins have high specificity for cardiac tissue. The cardiac troponins have a specific pattern of expression because they have different amino sequences encoded by different genes for skeletal and cardiac muscles. Cardiac TnI has an additional 31-amino acid residue compared to skeletal muscle. This protein is not expressed in normal, regenerating, or diseased skeletal muscle from human or animal origin (Bodor et al., 1995). Cardiac TnT has an additional 11-amino acid residue, but this protein was also found in regenerating rat skeletal muscle, during human fetal development, and in diseased human skeletal muscle (Anderson et al., 1991; Bodor et al., 1997; Saggin et al., 1990). In addition, cardiac TnT was also found in skeletal muscle specimens from patients with muscular dystrophy, polymyositis, and chronic renal disease (Bodor et al., 1997; McLaurin et al., 1997).

Neumann et al. (2019) evaluated high-sensitivity troponin (troponin I and T)’s ability to predict myocardial infarction and subsequent 30-day outcomes. The authors developed a risk assessment tool based on patients presenting to the emergency department with “symptoms suggestive of myocardial infarction”. Concentrations of troponin I or T were measured at presentation and after early or late serial sampling. Cutoffs were then determined to create cutoffs for risk assessment. Among the 22651 patients (9604 in derivation cohort, 13047 in validation cohort), the total prevalence of myocardial infarction was 15.3%. The authors found that “lower high-sensitivity troponin concentrations at presentation and smaller absolute changes during serial sampling were associated with a lower likelihood of myocardial infarction and a lower short-term risk of cardiovascular events” (Neumann et al., 2019).

Anand et al. (2019) evaluated the adoption rate of the universal definition of myocardial infarction and the corresponding recommendations. 1902 medical centers over 23 countries were surveyed, and the authors obtained answers regarding the primary biomarker, diagnostic thresholds, and clinical pathways used to identify myocardial infarction. The authors found that cardiac troponin was the primary biomarker used at 96% of surveyed sites, with 41% of these sites using high-sensitivity troponins. The sites using high-sensitivity assays were also more likely to use serial sampling (91% vs 78% using “contemporary” sensitivity troponin) and the 99th percentile diagnostic threshold (74% vs 66%). Use of creatine kinase-MB (CKMB) was “very limited” outside of Latin America (Anand et al., 2019).

In addition, other cardiac biomarkers, such as heart-type fatty acid binding protein (H-FABP) and copeptin, have been reported in the scientific literature. However, they are not commonly used in clinical settings (Jaffe & Morrow, 2021).

Boeddinghaus et al. (2020) compared the diagnostic accuracy of high-sensitivity cardiac troponin (hs-cTn) TriageTrue assay in patients with suspected myocardial infarction (MI) with other laboratory assays including f hs-cTnTElesys assay and hs-cTnI-Architect assay. 1,261 patients with symptoms suggestive of MI were enrolled in the study. The TriageTrue assay ruled out patients with troponin I concentration < 3 ng/l and classified these patients as low risk of MI and ruled in patients with a troponin I concentration > 60 ng/l. 178 out of the 1,261 patients enrolled in the study were diagnosed with MI based on troponin I levels of > 60 ng/l using the TriageTrue assay. TriageTrue troponin I concentrations were higher in patients with MI than in patients with other final diagnoses. Other diagnosis included unstable angina in 13 of 1,261 (9%), tachyarrhythmia, Takotsubo syndrome, heart failure, or myocarditis in 208 patients (17%), and noncardiac symptoms in 714 patients (57%). The AUC of the TriageTrue assay was 0.95, the hs-cTnT-Elecsys assay AUC was 0.93, and hs-cTnI-Architect assay AUC was 0.92. The TriageTrue algorithm allowed providers to make a triage decision after 1 hour in 401 of 545 patients. The efficacy for rule-out or rule-in was 43% for the TriageTrue, 25% in f hs-cTnTElesys, and 22% in hs-cTnI-Architect. Ruled-out patients had cumulative event rates of

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

0% at 30 days and 1.6% at 2 years. Overall, the authors conclude that “POC-hs-cTnI-TriageTrue assay provides high diagnostic accuracy in patients with suspected MI with a clinical performance that is at least comparable to that of best-validated central laboratory assays” (Boeddinghaus et al., 2020).

Heart-type fatty acid binding protein (H-FABP)

H-FABP, a small cytoplasmic protein present in cardiomyocytes, is believed to have a function in myocardial lipid homeostasis (Glatz & van der Vusse, 1990). Because of its small size, this protein appears in the blood after MI almost as early as myoglobin, but it has better specificity than myoglobin for cardiac tissue (Van Nieuwenhoven et al., 1995). Seino et al compared the use of H-FABP with rapid troponin in 371 patients with acute chest pain (Seino et al., 2003). Their study demonstrated that H-FABP had significantly higher sensitivity (89%) than troponin T (22%) and myoglobin (38%), but it has lower specificity (52%) than troponin (94%). Other studies were performed to compare H-FABP to troponins; however, they were unable to demonstrate superior results compared to troponins. H-FABP is not encouraged for assessment of MI as troponins are generally superior (Jaffe & Morrow, 2021).

In a prospective, cross-sectional study, Nguyen et al. (2020) studied the diagnostic utility of H-FABP in the early diagnosis of acute MI in comparison with troponin I and CK-MB. 216 patients enrolled in the study with 179 of those diagnosed with acute MI. H-FABP, CK-MB, and troponin I levels were compared. H-FABP reached its highest concentration in 6-12 hours after symptoms of chest pain, with a mean value of 169 ng/mL in acute MI patients. The cut-off value was 5.7 ng/ml with 90.5% sensitivity and 100% specificity. The combination of H-FABP, CK-MB and troponin I together had the highest sensitivity of 97.2%. The AUC of H-FABP was observed to be 0.99, which was higher than CK-MB (0.92) and troponin I (0.86). The authors conclude that “H-FABP can be used as a reliable diagnostic cardiac biomarker in the early detection of AMI for patients who came to the emergency room within 12h of onset of chest pain” (Nguyen et al., 2020).

Copeptin

Copeptin is the 39 amino acid C-terminal fragment cleaved from pro-arginine vasopressin (AVP). After MI, copeptin levels increase rapidly and decline over the next two to five days (Khan et al., 2007). In the Copeptin Helps in the Early Detection of Patients With Acute Myocardial Infarction (CHOPIN) 16-site study involving 1,967 patients presenting within 6 hours of pain onset, copeptin was shown to have a potential value in ruling out MI with a negative predictive value greater than 99% when combined with TnI measurements (Maisel et al., 2013). The Advantageous Predictors of Acute Coronary Syndrome Evaluation (APACE) multicenter study, involving 1,439 patients presenting with MI symptoms, demonstrated no benefit in using copeptin as an early rule-out cardiac biomarker for MI (Hillinger et al., 2015). Copeptin is not encouraged for assessment of MI as troponins are generally superior (Jaffe & Morrow, 2021).

ST2

ST2 is part of the interleukin-1 receptor family with two isoforms, soluble ST2 (sST2) and ST2L. ST2 is the receptor of the IL-33 cytokine that can be secreted by living cells in response to cellular stress and mechanical strain. IL-33 binds the receptor complex of ST2L and IL-1R accessory protein and reduces myocardial fibrosis, prevents cardiomyocyte hypertrophy, reduces apoptosis, and improves myocardial function. The cardioprotective effects of IL-33 are specifically through the ST2L receptor. However, sST2 may also bind IL-33, blocking the interaction between IL-33/ST2L. This eliminates the cardioprotective effects of the IL-33/ST2L interaction (Pascual-Figal & Januzzi, 2015). Experimentally, this leads to cardiac hypertrophy, fibrosis, and ventricular dysfunction (Januzzi et al., 2015).

One of the main proprietary tests used to assess ST2 levels is the Presage Assay by Critical Diagnostics. This assay was approved by the FDA on December 9, 2011. According to the FDA, “The Critical Diagnostics Presage® ST2 Assay kit is an in vitro diagnostic device that quantitatively measures ST2 in serum or plasma by enzyme-linked immunosorbent assay (ELISA) in a microtiter plate format. The Presage® ST2 Assay is indicated to be used in conjunction with clinical evaluation as an aid in

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

assessing the prognosis of patients diagnosed with chronic heart failure”. The manufacturer claims a measuring range of 3.1 ng/mL of soluble ST2 to 200 ng/mL, and the data based on 1100 samples supports this claim. These 1100 samples found coefficient of variation of under 5%, a linear curve, and a $r^2=0.99$ (FDA, 2011).

Proprietary Testing

Proprietary tests for various biomarkers are available in several clinical settings. Platforms including Roche’s “CARDIAC Trop T Sensitive test” and ResponseBio’s battery of cardiac tests emphasize their speed (on the scale of minutes) and versatility (ResponseBio, 2020; Roche, 2020).

No single diagnostic test for HF exists because it is largely a clinical diagnosis based on a careful history and physical examination. However, biomarkers of cardiovascular diseases have been developed for diagnosis and prognosis, and the use of several biomarkers is now considered the standard of care. ST2 is a marker of cardiomyocyte stress and fibrosis that adds additional value to natriuretic peptides, resulting in a risk stratification of patients with a wide spectrum of cardiovascular diseases (Bayes-Genis et al., 2015).

Clinical Utility and Validity

Jeong et al. (2020) studied the diagnostic value of copeptin for early diagnosis of acute MI in comparison with troponin I and CK-MB. 271 patients complaining of chest pain within 6 hours of onset were studied within the emergency department. The diagnostic performance of copeptin, troponin I, and CK-MB was compared by assessing the AUC and ROC curve analysis. After comparing AUC, copeptin had a significantly better diagnosis value than troponin I in patients with chest pain within two hours of onset. In addition, troponin I and copeptin together had better diagnostic performance than CK-MB and troponin I combination. Overall, the authors conclude that “the combination of troponin I and copeptin improves AMI diagnostic performance in patients with early-onset chest pain in an ED setting” (Jeong et al., 2020).

conducted a multi-center prospective study to evaluate whether plasma ST2 levels predict adverse outcomes in 1,141 chronic heart failure outpatients. Patients in the highest ST2 tertile (ST2 > 36.3 ng/mL) had a “markedly increased” risk (hazard ratio 3.2) of adverse outcomes compared to the lowest tertile ≤ 22.3 ng/mL). The investigators concluded that “ST2 is a potent marker of risk in chronic heart failure and when used in combination with NT-proBNP offers moderate improvement in assessing prognosis beyond clinical risk scores” (Ky et al., 2011).

Wang et al. (2012) studied the prognostic value of three novel biomarkers induced by cardiovascular stress. The investigators measured sST2, growth differentiation factor-15, and high-sensitivity troponin I in 3,428 participants in the Framingham Heart Study. Multivariable-adjusted proportional hazards models were performed to assess the individual and combined ability of the biomarkers to predict adverse outcomes. The three new biomarkers were associated with death, major cardiovascular events, and heart failure, but not with coronary events. The investigators concluded that the findings demonstrated the prognostic value of the newer biomarkers in apparently healthy individuals (Wang et al., 2012).

Wang et al. (2018) investigated the possibility of using sST2 as a biomarker to distinguish between acute aortic dissection and other causes of acute chest pain. Using an R&D Systems assay to measure plasma concentrations of sST2 in 1360 patients with a cutoff of 34.6 ng/mL, the researchers found that “sST2 had a sensitivity of 99.1%, specificity of 84.9%, positive predictive value of 68.7%, negative predictive value of 99.7%, positive likelihood ratio of 6.6, and negative likelihood ratio of 0.01.” Additionally, within 24 hours of symptom onset, sST2 levels were higher in those with acute aortic dissection in comparison to those with acute myocardial infarction or pulmonary embolism. sST2 was also superior in overall diagnostic performance to D-dimer and troponin I using the area under receiver operating characteristic curves.

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

Januzzi et al. (2013) conducted a retrospective study to assess sST2 as a prognostic marker after orthotopic heart transplantation (OHT) and as a test to predict acute cellular rejection. sST2 concentrations were measured in 241 patients following OHT. Elevated sST2 was associated with cellular rejection, with highest rates of cellular rejection in the 4th sST2 quartile. No significant association between sST2 and antibody-mediated rejection or allograft vasculopathy was found. A sST2 level of ≥ 30 ng/mL was found to independently predict death over the 7-year follow-up with a hazard ratio of 2.1. The investigators concluded that sST2 levels are associated with the presence of cellular rejection and predict long-term mortality following OHT (Januzzi et al., 2013).

Boman et al. (2018) assessed the prognostic value of ST2 on cardiovascular mortality. 159 patients were evaluated, but ST2 was not found to be significantly associated with cardiovascular mortality or all-cause mortality. Furthermore, no significant interaction of ST2 and N-terminal pro-hormone of brain natriuretic peptide /N-terminal pro-B-type natriuretic peptide was found (Boman et al., 2018).

Dimitropoulos et al. (2020) investigated the association of soluble suppression of tumorigenesis-2 (sST2) with endothelial function in patients with ischemic heart failure. A total of 143 patients with “table HF of ischemic etiology and reduced left ventricular ejection fraction (LVEF)” were included along with 77 controls. The authors found an increased level of sST2 in HF patients compared to controls (15.8 ng/mL compared to 12.5 ng/mL). Within the HF group, there was no association of LVEF with sST2. Overall, sST2 levels were found to be increased and associated with functional capacity in “patients with chronic HF of ischemic etiology”. Finally, the authors found an inverse association between flow-mediated dilation and sST2 levels, which the authors stated “highlight[ed] the interplay between the dysfunctional endothelium and HF pathophysiologic mechanisms” (Dimitropoulos et al., 2020).

Hou et al. (2020) aimed to investigate the association between sST2 levels and clinical outcomes of high-risk heart failure. The primary endpoint was defined as all-cause mortality. A total of 150 patients were included; all-cause mortality occurred in 16 of the patients over the course follow-up. The authors found that all-cause mortality increased significantly above 34.98846 ng/mL by a factor of 16% to 5.33%. After adjusting the model for certain co-factors (age, gender, et al.), and after adding NT-proBNP, “the risk of all-cause death was increased by 2.5% and 1.9%, respectively, per ng/ml of sST2”. The authors identified the best sST2 cut-off for predicting all-cause mortality to be 43.42671 ng/ml, with an area under the curve of 0.72, sensitivity of 0.69, and specificity of 0.69. Risk of all-cause mortality was found to be 21.2% above this cutoff and 5.1% below it, with a corresponding hazard ratio of 3.30. The authors concluded that “Patients with sST2 levels more than 43.42671 ng/ml even after ICD implantation should therefore be monitored carefully” (Hou et al., 2020).

Guidelines and Recommendations

2018 ESC/ACC/AHA/WHF Fourth Universal Definition of Myocardial Infarction

Both cTnI and cTnT are recommended for evaluation of myocardial injury, and high sensitivity cTn assays are recommended for routine clinical use. An acute MI is designated when a rising/falling pattern is seen with cTn levels and if there is at least one measurement greater than the 99th percentile of the upper reference limit (URL) (Jaffe et al., 2018).

CKMB is considered less sensitive and specific than either troponin. However, in the absence of a cTn assay, CK-MB is considered the best alternative. A measurement of CK-MB above the 99th percentile of the URL should be “designated as the decision level for the diagnosis of MI”. Sex-specific CK-MB values should be used (Jaffe et al., 2018).

In the 2019 AHA guideline discussing the “Contemporary Diagnosis and Management of Patients With Myocardial Infarction in the Absence of Obstructive Coronary Artery Disease [MINOCA]”, the AHA notes that the diagnostic criteria of MINOCA follows the “Fourth Universal Definition of Myocardial Infarction” above, specifically the rise or fall of cardiac troponin levels with at least one value above the

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS

– G2150 “Notification”

99th percentile of the reference limit. The guideline considers this definition “fundamental” to identifying and defining MINOCA (Tamis-Holland Jacqueline et al., 2019).

2014 AHA/ACC Guideline for the Management of Patients with Non-ST-Elevation Acute Coronary Syndromes (NSTE-ACS)

The American College of Cardiology (ACC) and the American Heart Association (AHA) have developed clinical practice guidelines to provide recommendations applicable to patients with or at risk of developing cardiovascular disease and to provide guidance to clinicians on optimal management of patients with NSTE-ACS. In their comprehensive document, the AHA/ACC panel has provided recommendations for initial evaluation and management of patients presenting with ACS symptoms, for the early hospital care, myocardial revascularization, late hospital care, hospital discharge and posthospital discharge care, special patient groups and quality of care and outcomes for ACS. The Task Force recommended to stratify patients with suspected ACS based on the likelihood of ACS and those with high-risk features should be referred immediately to the emergency department (ED). They have provided specific recommendations for the use of cardiac biomarkers in the diagnosis and prognosis of MI. They specifically recommended using troponin (troponin I or T when contemporary assay is used) for the diagnosis of MI. According to AHA/ACC guidelines, the cardiac troponin is recommended and should be measured at presentation and 3 to 6 hours after symptom onset in all patients who present with ACS symptoms. The panelists recommended identifying rising and/or falling pattern of troponin. In addition, they recommended measuring troponin levels beyond 6 hours after symptom onset in patients with normal troponins on serial examination when ECG changes and/or clinical presentation suggests ACS. If the onset of symptoms is not clearly identified, they recommended using the time of presentation as the time of onset for measuring troponin. The AHA/ACC guideline clearly highlighted that CKMB, or myoglobin should not be used for the diagnosis of ACS. All recommendations for the use of cardiac biomarkers in the diagnosis of MI were level A evidence.

The AHA/ACC guideline considered all recommendations in the use of cardiac biomarkers for ACS prognosis as level of evidence B. They considered the presence and magnitude of troponin elevations useful for short- and long-term prognosis. The re-measurement of troponin once on day 3 or day 4 in patients with MI was considered reasonable to estimate the infarct size and dynamics of necrosis. Finally, they considered the use of B-type natriuretic peptide to be reasonable for additional prognostic information.

The recommendations for the use of cardiac biomarkers in the diagnosis and prognosis of MI was well summarized in Table from 2014 AHA/ACC guidelines p.2655 (Amsterdam et al., 2014):

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

TABLE 5 Summary of Recommendations for Cardiac Biomarkers and the Universal Definition of MI

Recommendations	COR	LOE
Diagnosis		
Measure cardiac-specific troponin (troponin I or T) at presentation and 3–6 h after symptom onset in all patients with suspected ACS to identify pattern of values	I	A
Obtain additional troponin levels beyond 6 h in patients with initial normal serial troponins with electrocardiographic changes and/or intermediate/high risk clinical features	I	A
Consider time of presentation the time of onset with ambiguous symptom onset for assessing troponin values	I	A
With contemporary troponin assays, CK-MB and myoglobin are not useful for diagnosis of ACS	III: No Benefit	A
Prognosis		
Troponin elevations are useful for short- and long-term prognosis	I	B
Remeasurement of troponin value once on d 3 or 4 in patients with MI may be reasonable as an index of infarct size and dynamics of necrosis	IIb	B
BNP may be reasonable for additional prognostic information	IIb	B

ACS indicates acute coronary syndromes; BNP, B-type natriuretic peptide; CK-MB, creatine kinase myocardial isoenzyme; COR, Class of Recommendation; LOE, Level of Evidence; and MI, myocardial infarction.

2013 (published 2014) Society for Cardiovascular Angiography and Interventions (SCAI)

In their expert consensus document titled “Consideration of a New Definition of Clinically Relevant Myocardial Infarction After Coronary Revascularization,” the SCAI expert panel introduced a new definition of clinically relevant MI after coronary revascularization percutaneous coronary intervention (PCI) or coronary artery bypass grafting (CABG). In their definition of clinically relevant MI after both PCI and CABG procedures, authors gave recommendations according to 3 different types of clinical presentation. In the first case, when patient has a normal CKMB baseline: “The peak CK-MB measured within 48 hours of the procedure rises to $\geq 10x$ the local laboratory ULN, or to $\geq 5x$ ULN with new pathologic Q-waves in ≥ 2 contiguous leads or new persistent LBBB, OR in the absence of CK-MB measurements and a normal baseline cTn, a cTn (I or T) level measured within 48 hours of the PCI rises to $\geq 70x$ the local laboratory ULN, or $\geq 35x$ ULN with new pathologic Q-waves in ≥ 2 contiguous leads or new persistent LBBB”. In the case when patients have elevated baseline CKMB (or cTn) with stable or falling biomarkers levels, they issued the following recommendation: “The CK-MB (or cTn) rises by an absolute increment equal to those levels recommended above from the most recent pre-procedure level”. And, in patients with elevated CKMB (or cTn), but without stable or falling biomarkers level, the recommendation was: “The CK-MB (or cTn) rises by an absolute increment equal to those levels recommended above plus new ST-segment elevation or depression plus signs consistent with a clinically relevant MI, such as new onset or worsening heart failure or sustained hypotension”. The authors have expressed preference to use CKMB instead of cTn, but they have included cTn in their definition if CKMB was not available (Moussa et al., 2013).

2015 AHA Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care

In their review of previously issued guidelines, the expert panel introduced new recommendations for diagnostic interventions in ACS regarding cardiac biomarkers. They still recommended to use Troponin in following situations: “We recommend against using hs-cTnT and cTnI alone measured at 0 and 2 hours (without performing clinical risk stratification) to identify patients at low risk for ACS (Class III: Harm, LOE B-NR). We recommend that hs-cTnI measurements that are less than the 99th percentile, measured at 0 and 2 hours, may be used together with low-risk stratification (TIMI score of 0 or 1 or low risk per Vancouver rule) to predict a less than 1% chance of 30-day MACE (Class IIa, LOE B-NR). We recommend that negative cTnI or cTnT measurements at 0 and between 3 and 6 hours may be used together with very low-risk stratification (TIMI score of 0, low-risk score per Vancouver rule, North American Chest Pain score of 0 and age less than 50 years, or low-risk HEART score) to predict a less than 1% chance of 30-day MACE (Class IIa, LOE B-NR)”. They did not express a preference in cardiac biomarkers to use, nor did they give any recommendations regarding CKMB (O’Connor Robert et al., 2015).

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

American College of Cardiology/American Heart Association/Heart Failure Society of America (ACC/AHA/HFSA)

In 2017, the ACC, AHA, and HFSA jointly updated evidence-based guidelines for the management of heart failure (HF). The ACC/AHA/HFSA stated that “Biomarkers of myocardial fibrosis, soluble ST2 receptor, and galectin-3 are predictive of hospitalization and death and may provide incremental prognostic value over natriuretic peptide levels in patients with HF.” The guidelines also note that “in patients with chronic HF, clinically available tests such as biomarkers of myocardial injury or fibrosis may be considered for additive risk stratification in patients” (Yancy et al., 2017).

American Heart Association (AHA)

The AHA notes sST2 as an “emerging” biomarker that supports diagnosis of HF with preserved ejection fraction, a biomarker that may predict mortality and HF events, and a biomarker that correlates with left ventricular end-diastolic pressure. The AHA states that sST2 has numerous advantages as a biomarker, namely its concentration being unaffected by BMI, age, or renal function. SST2 is stated to correlate with HF prognosis as well. Overall, AHA states that out of the newer biomarkers (SST2, ST2, Gal-3, and GDF-15), “most appeal is driven by sST2” (Chow et al., 2017).

A Scientific Statement published in 2019 also considered ST2 as the most “promising clinically”, but also mentioned the limitations in consistency and utility in most inflammatory mediators. The Statement notes several clinical studies focusing on sST2 that are in progress as of March 24, 2020 (Cresci et al., 2019).

European Society of Cardiology (ESC)

The ESC notes measurement of cardiac troponins as “mandatory” in all patients with suspected non-ST-elevation acute coronary syndromes. The guidelines assert that cardiac troponins are more sensitive and specific biomarkers of cardiomyocyte injury than CK, CKMB, and myoglobin. However, if troponin measurement is not possible, measurement of copeptin is recommended.

The ESC also acknowledges the natriuretic peptides (B-type natriuretic peptide, N-terminal pro-B-type natriuretic peptide and midregional pro-A-type natriuretic peptide) as providing useful prognostic information along with the troponins. The ESC mentions other biomarkers such as midregional pro-adrenomedullin, growth differentiation factor 15 and copeptin, but they cannot recommend them at this time as their added value in risk assessment seems “marginal” (Gencer et al., 2016).

The 2019 ESC guidelines focusing on chronic coronary syndromes states that for “clinical suspicion of coronary artery disease instability...management should follow the Guidelines for ACS without persistent ST-segment elevation”, which is discussed above (Knuuti et al., 2019).

The 2020 ESC guidelines focus on diagnosis of acute coronary syndrome. Regarding MI, they recommend that “the routine use of copeptin as an additional biomarker for the early rule-out of MI should be considered where hs-cTn assays are not available.” In addition, “CK-MB shows a more rapid decline after MI and may provide added value for detection of early reinfarction” (Collet et al., 2021).

In their 2016 guidelines on acute and chronic heart failure, the ESC states that “although there is extensive research on biomarkers in HF (e.g. ST2, galectin 3, copeptin, adrenomedullin), there is no definite evidence to recommend them for clinical practice” (Ponikowski et al., 2016).

However, in a 2021 update, ESC (with special contribution from the Heart Failure Association) lists the key elements for HF and CMP diagnostic workups, and among the laboratory exams recommended under Table 26 for the “Initial diagnostic assessment in patients with suspected cardiomyopathy”, they include the use of ST2:

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

“**Laboratory exams** including cardiac and muscular enzymes, liver and renal function, haemoglobin, white blood cell count (including differential white blood cell count to detect eosinophilia), natriuretic peptides, thyroid function tests, iron status, and markers of systemic auto-immune disease (hsCRP, anti-nuclear antibodies, soluble IL-2 receptor)” (McDonagh et al., 2021)

Heart Failure Association of the European Society of Cardiology

The Heart Failure Association of the European Society of Cardiology published a position statement on Advanced Heart Failure which states: “Post-transplant patients should undergo a pre-defined regimen of graft biopsies, titration of immunosuppressive and other therapies, rejection monitoring, assessment for infections, transplant coronary artery disease and/or cardiac allograft vasculopathy, immunosuppression side effects, and other potential complications including neoplasia, and co-morbidities that require comprehensive treatment.” However, the guideline does not mention sST2 regarding prognosis of post-transplant patients (Crespo-Leiro et al., 2018).

ACC/AATS/AHA/ASE/ASNC/SCAI/SCCT/STS 2016 Appropriate Use Criteria for Coronary Revascularization in Patients With Acute Coronary Syndromes Guidelines

In 2016 The American College of Cardiology (ACC), Society for Cardiovascular Angiography and Interventions (SCAI), Society of Thoracic Surgeons (STS), and American Association for Thoracic Surgery (AATS), along with key specialty and subspecialty societies created an Appropriate Use Task Force with the mission to revise the appropriate use criteria (AUC) for coronary revascularization. They have used clinical scenarios to mimic patient presentations seen in everyday clinical practice and included information on symptom status, presence of clinical instability or ongoing ischemic symptoms and other characteristics. They follow 2014 AHA/ACC recommendations for the use of cardiac biomarkers (Amsterdam et al., 2014).

American Society for Clinical Pathology (ASCP)

The ASCP recommends against testing CK-MB or myoglobin to diagnose an acute myocardial infarction. Instead, they recommend testing either troponin I or T. They also assert that both troponins are specific to cardiac injury and that there is much support for relying solely on troponin (ASCP, 2015).

National Institute for Health and Care Excellence (NICE)

NICE recommends diagnosis of MI using the “detection of rise and/or fall of cardiac biomarkers values [preferably cardiac troponin (cTn)] with at least one value above the 99th percentile of the upper reference limit and at least one of the following:

- symptoms of ischaemia
- new or presumed new significant ST-segment-T wave (ST-T) changes or new left bundle branch block (LBBB)
- development of pathological Q waves in the ECG
- imaging evidence of new loss of viable myocardium or new regional wall motion abnormality
- identification of an intracoronary thrombus by angiography” (NICE, 2016).

Currently, the 2018 NICE recommendations on chronic heart failure do not mention the usage of ST2 as a marker for diagnosing chronic heart failure. Instead, they recommend to “measure N-terminal pro-B-type natriuretic peptide (NT-proBNP) in people with suspected heart failure” (NICE, 2018).

In 2020, NICE released recommendations on the use of high sensitivity troponin tests to help rule out NSTEMI earlier in those presenting to an emergency department with chest pain and suspected acute coronary syndrome. NICE recommends the use of the following assays: Access High-Sensitivity Troponin I Assay, ADVIA Centaur High-Sensitivity Cardiac Troponin-I Assay, Alinity High Sensitive Troponin-I assay, ARCHITECT STAT High Sensitive Troponin-I assay, Atellica IM High-Sensitivity Cardiac Troponin I Assay, Dimension Vista High-Sensitivity Cardiac Troponin I Assay, Dimension EXL High-Sensitivity Cardiac Troponin I Assay, Elecsys Troponin T-high sensitive assay, Elecsys Troponin T-high sensitive STAT assay, VIDAS High sensitive

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

Troponin I assay, and VITROS High Sensitivity Troponin I Assay. NICE mentions that although the “TriageTrue test has the potential to be cost effective, its diagnostic accuracy when used on whole blood is uncertain” (NICE, 2020). Regarding use of these assays, NICE recommends using a threshold at or near the limit of detection, which varies depending on the assay used. If this sample is positive, it should not be used to rule in NSTEMI. If taking multiple samples, take a sample at initial assessment followed by a second sample taken 30 minutes to 3 hours after. Use 99th percentile thresholds or thresholds at or near the limit of detection of the assay (NICE, 2020).

Canadian Cardiovascular Society (CCS)

The CCS notes sST2 as a potential prognostic biomarker and states that “might be superior” to galectin-3. However, the CCS also remarks that it is “unclear” if using sST2 in HF to modify therapies improve clinical outcomes (Ezekowitz et al., 2017).

State and Federal Regulations, as applicable

Food and Drug Administration (FDA)

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

Billing/Coding/Physician Documentation Information

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at www.bcbsnc.com. They are listed in the Category Search on the Medical Policy search page.

Applicable service codes: 82550, 82552, 82553, 82554, 82725, 83006, 83615, 83625, 83874, 84450, 84484, 84512, 84588, 84999, 86140

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful, but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

Scientific Background and Reference Sources

Amsterdam, E. A., Wenger, N. K., Brindis, R. G., Casey, D. E., Ganiats, T. G., Holmes, D. R., Jaffe, A. S., Jneid, H., Kelly, R. F., Kontos, M. C., Levine, G. N., Liebson, P. R., Mukherjee, D., Peterson, E. D., Sabatine, M. S., Smalling, R. W., & Zieman, S. J. (2014). 2014 AHA/ACC Guideline for the Management of Patients With Non–ST-Elevation Acute Coronary Syndromes. *A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines*, 130(25), e344-e426. <https://doi.org/10.1161/cir.000000000000134>

Anand, A., Shah, A. S. V., Beshiri, A., Jaffe, A. S., & Mills, N. L. (2019). Global Adoption of High-Sensitivity Cardiac Troponins and the Universal Definition of Myocardial Infarction. *Clin Chem*, 65(3), 484-489. <https://doi.org/10.1373/clinchem.2018.298059>

Anderson, P. A., Malouf, N. N., Oakeley, A. E., Pagani, E. D., & Allen, P. D. (1991). Troponin T isoform expression in humans. A comparison among normal and failing adult heart, fetal heart, and adult and fetal skeletal muscle. *Circ Res*, 69(5), 1226-1233. <https://doi.org/10.1161/01.res.69.5.1226>

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

- ASCP. (2015, February 3). *Don't test for myoglobin or CK-MB in the diagnosis of acute myocardial infarction (AMI). Instead, use troponin I or T.* <https://www.choosingwisely.org/clinician-lists/american-society-clinical-pathology-myoglobin-to-diagnose-acute-myocardial-infarction/>
- Bayes-Genis, A., Zhang, Y., & Ky, B. (2015). ST2 and patient prognosis in chronic heart failure. *Am J Cardiol*, 115(7 Suppl), 64b-69b. <https://doi.org/10.1016/j.amjcard.2015.01.043>
- Bessman, S. P., & Carpenter, C. L. (1985). The creatine-creatine phosphate energy shuttle. *Annu Rev Biochem*, 54, 831-862. <https://doi.org/10.1146/annurev.bi.54.070185.004151>
- Bodor, G. S., Porterfield, D., Voss, E. M., Smith, S., & Apple, F. S. (1995). Cardiac troponin-I is not expressed in fetal and healthy or diseased adult human skeletal muscle tissue. *Clin Chem*, 41(12 Pt 1), 1710-1715. <https://pubmed.ncbi.nlm.nih.gov/7497610/>
- Bodor, G. S., Survant, L., Voss, E. M., Smith, S., Porterfield, D., & Apple, F. S. (1997). Cardiac troponin T composition in normal and regenerating human skeletal muscle. *Clin Chem*, 43(3), 476-484. <https://doi.org/10.1093/clinchem/43.3.476>
- Boeddinghaus, J., Nestelberger, T., Koechlin, L., Wussler, D., Lopez-Ayala, P., Walter, J. E., Troester, V., Ratmann, P. D., Seidel, F., Zimmermann, T., Badertscher, P., Wildi, K., Giménez, M. R., Potlukova, E., Strebel, I., Freese, M., Miró, Ò., Martin-Sanchez, F. J., Kawecki, D., . . . Geigy, N. (2020). Early Diagnosis of Myocardial Infarction With Point-of-Care High-Sensitivity Cardiac Troponin I. *Journal of the American College of Cardiology*, 75(10), 1111-1124. <https://doi.org/doi:10.1016/j.jacc.2019.12.065>
- Boman, K., Thormark Frost, F., Bergman, A. R., & Olofsson, M. (2018). NTproBNP and ST2 as predictors for all-cause and cardiovascular mortality in elderly patients with symptoms suggestive for heart failure. *Biomarkers*, 23(4), 373-379. <https://doi.org/10.1080/1354750x.2018.1431692>
- Chow, S. L., Maisel, A. S., Anand, I., Bozkurt, B., de Boer Rudolf, A., Felker, G. M., Fonarow, G. C., Greenberg, B., Januzzi, J. L., Kiernan, M. S., Liu, P. P., Wang, T. J., Yancy, C. W., & Zile, M. R. (2017). Role of Biomarkers for the Prevention, Assessment, and Management of Heart Failure: A Scientific Statement From the American Heart Association. *Circulation*, 135(22), e1054-e1091. <https://doi.org/10.1161/CIR.0000000000000490>
- Collet, J. P., Thiele, H., Barbato, E., Barthélémy, O., Bauersachs, J., Bhatt, D. L., Dendale, P., Dorobantu, M., Edvardsen, T., Folliguet, T., Gale, C. P., Gilard, M., Jobs, A., Jüni, P., Lambrinou, E., Lewis, B. S., Mehilli, J., Meliga, E., Merkely, B., . . . Siontis, G. C. M. (2021). 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J*, 42(14), 1289-1367. <https://doi.org/10.1093/eurheartj/ehaa575>
- Colucci, W. (2021, January 27). *Prognosis of heart failure.* <https://www.uptodate.com/contents/prognosis-of-heart-failure>
- Colucci, W. (2022, November 9, 2022). *Overview of the therapy of heart failure with reduced ejection fraction.* <https://www.uptodate.com/contents/overview-of-the-management-of-heart-failure-with-reduced-ejection-fraction-in-adults>
- Colucci, W., & Dunlay, S. (2022, August 10, 2022). *Clinical manifestations and diagnosis of advanced heart failure.* <https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-advanced-heart-failure>

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

Cresci, S., Pereira Naveen, L., Ahmad, F., Byku, M., de las Fuentes, L., Lanfear David, E., Reilly Carolyn, M., Owens Anjali, T., Wolf Matthew, J., & null, n. (2019). Heart Failure in the Era of Precision Medicine: A Scientific Statement From the American Heart Association. *Circulation: Genomic and Precision Medicine*, 12(10), e000058.

<https://doi.org/10.1161/HCG.0000000000000058>

Crespo-Leiro, M. G., Metra, M., Lund, L. H., Milicic, D., Costanzo, M. R., Filippatos, G., Gustafsson, F., Tsui, S., Barge-Caballero, E., De Jonge, N., Frigerio, M., Hamdan, R., Hasin, T., Hulsmann, M., Nalbantgil, S., Potena, L., Bauersachs, J., McDonagh, T., Seferovic, P., & Ruschitzka, F. (2018). Advanced heart failure: a position statement of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail*.

<https://doi.org/10.1002/ehf.1236>

Danese, E., & Montagnana, M. (2016). An historical approach to the diagnostic biomarkers of acute coronary syndrome. *Ann Transl Med*, 4(10), 194. <https://doi.org/10.21037/atm.2016.05.19>

Dillon, M. C., Calbreath, D. F., Dixon, A. M., Rivin, B. E., Roark, S. F., Ideker, R. E., & Wagner, G. S. (1982). Diagnostic problem in acute myocardial infarction: CK-MB in the absence of abnormally elevated total creatine kinase levels. *Arch Intern Med*, 142(1), 33-38.

<https://doi.org/10.1001/archinte.1982.00340140035009>

Dimitropoulos, S., Mystakidi, V. C., Oikonomou, E., Siasos, G., Tsigkou, V., Athanasiou, D., Gouliopoulos, N., Bletsas, E., Kalampogias, A., Charalambous, G., Tsioufis, C., Vavuranakis, M., & Tousoulis, D. (2020). Association of Soluble Suppression of Tumorigenesis-2 (ST2) with Endothelial Function in Patients with Ischemic Heart Failure. *Int J Mol Sci*, 21(24).

<https://doi.org/10.3390/ijms21249385>

Eggers, K. M., Oldgren, J., Nordenskjold, A., & Lindahl, B. (2004). Diagnostic value of serial measurement of cardiac markers in patients with chest pain: limited value of adding myoglobin to troponin I for exclusion of myocardial infarction. *Am Heart J*, 148(4), 574-581.

<https://doi.org/10.1016/j.ahj.2004.04.030>

Engel, G., & Rockson, S. G. (2020). Feasibility and Reliability of Rapid Diagnosis of Myocardial Infarction. *The American Journal of the Medical Sciences*, 359(2), 73-78.

<https://doi.org/10.1016/j.amjms.2019.12.012>

Ezekowitz, J., O'Meara, E., McDonald, M., Abrams, H., Chan, M., Ducharme, A., Giannetti, N., Grzeslo, A., Hamilton, P., Heckman, G., Howlett, J., Koshman, S., Lepage, S., McKelvie, R., Moe, G., Rajda, M., Swiggum, E., Virani, S., Zieroth, S., . . . Sussex, B. (2017). 2017 *Comprehensive Update of the Canadian Cardiovascular Society Guidelines for the Management of Heart Failure*. [https://www.onlinecjc.ca/article/S0828-282X\(17\)30973-X/pdf](https://www.onlinecjc.ca/article/S0828-282X(17)30973-X/pdf)

FDA. (2011). *SUBSTANTIAL EQUIVALENCE DETERMINATION*

https://www.accessdata.fda.gov/cdrh_docs/reviews/K111452.pdf

Gencer, B., Brotons, C., Mueller, C., Mukherjee, D., Chew, D. P., Andreotti, F., Hasenfuss, G., Collet, J.-P., Bax, J. J., Mehilli, J., Kjeldsen, K., Valgimigli, M., Borger, M. A., Lancellotti, P., Storey, R. F., Windecker, S., Landmesser, U., Patrono, C., Roffi, M., & Group, E. S. C. S. D. (2016). 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *European Heart Journal*, 37(3), 267-315.

<https://doi.org/10.1093/eurheartj/ehv320>

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

- Glatz, J. F., & van der Vusse, G. J. (1990). Cellular fatty acid-binding proteins: current concepts and future directions. *Mol Cell Biochem*, 98(1-2), 237-251. <https://doi.org/10.1007/bf00231390>
- Greaser, M. L., & Gergely, J. (1971). Reconstitution of troponin activity from three protein components. *J Biol Chem*, 246(13), 4226-4233. [https://doi.org/10.1016/S0021-9258\(18\)62075-7](https://doi.org/10.1016/S0021-9258(18)62075-7)
- Heller, G. V., Blaustein, A. S., & Wei, J. Y. (1983). Implications of increased myocardial isoenzyme level in the presence of normal serum creatine kinase activity. *Am J Cardiol*, 51(1), 24-27. [https://doi.org/10.1016/s0002-9149\(83\)80006-x](https://doi.org/10.1016/s0002-9149(83)80006-x)
- Hillinger, P., Twerenbold, R., Jaeger, C., Wildi, K., Reichlin, T., Gimenez, M. R., Engels, U., Miró, O., Boeddinghaus, J., Puelacher, C., Nestelberger, T., Röthlisberger, M., Ernst, S., Rentsch, K., & Mueller, C. (2015). Optimizing Early Rule-Out Strategies for Acute Myocardial Infarction: Utility of 1-Hour Copeptin. *Clin Chem*, 61(12), 1466-1474. <https://doi.org/10.1373/clinchem.2015.242743>
- Hou, Z. W., Yu, H. B., Liang, Y. C., Gao, Y., Xu, G. Q., Wu, M., Mei, Z., Wang, Z. L., Li, Z. G., Li, Y. Y., Song, H. X., Li, J. Y., & Han, Y. L. (2020). Circulating Soluble ST2 Predicts All-Cause Mortality in Severe Heart Failure Patients with an Implantable Cardioverter Defibrillator. *Cardiol Res Pract*, 2020, 4375651. <https://doi.org/10.1155/2020/4375651>
- Jaffe, A. S., Chaitman, B. R., Morrow, D. A., Bax, J. J., White, H. D., Alpert, J. S., Thygesen, K., & Group, E. S. C. S. D. (2018). Fourth universal definition of myocardial infarction (2018). *European Heart Journal*, 40(3), 237-269. <https://doi.org/10.1093/eurheartj/ehy462>
- Jaffe, A. S., & Morrow, D. A. (2021, February 15). *Biomarkers of cardiac injury other than troponin*. Wolters Kluwer. <https://www.uptodate.com/contents/biomarkers-of-cardiac-injury-other-than-troponin>
- Januzzi, J. L., Horne, B. D., Moore, S. A., Galenko, O., Snow, G. L., Brunisholz, K. D., Muhlestein, J. B., Alharethi, R., Carlquist, J. F., Budge, D., Rasmussen, K., & Kfoury, A. G. (2013). Interleukin receptor family member ST2 concentrations in patients following heart transplantation. *Biomarkers*, 18(3), 250-256. <https://doi.org/10.3109/1354750x.2013.773081>
- Januzzi, J. L., Mebazaa, A., & Di Somma, S. (2015). ST2 and prognosis in acutely decompensated heart failure: the International ST2 Consensus Panel. *Am J Cardiol*, 115(7 Suppl), 26b-31b. <https://doi.org/10.1016/j.amjcard.2015.01.037>
- Jeong, J. H., Seo, Y. H., Ahn, J. Y., Kim, K. H., Seo, J. Y., Chun, K. Y., Lim, Y. S., & Park, P. W. (2020). Performance of Copeptin for Early Diagnosis of Acute Myocardial Infarction in an Emergency Department Setting. *Ann Lab Med*, 40(1), 7-14. <https://doi.org/10.3343/alm.2020.40.1.7>
- Kavsak, P. A., MacRae, A. R., Newman, A. M., Lustig, V., Palomaki, G. E., Ko, D. T., Tu, J. V., & Jaffe, A. S. (2007). Effects of contemporary troponin assay sensitivity on the utility of the early markers myoglobin and CKMB isoforms in evaluating patients with possible acute myocardial infarction. *Clin Chim Acta*, 380(1-2), 213-216. <https://doi.org/10.1016/j.cca.2007.01.001>
- Khan, S. Q., Dhillon, O. S., O'Brien, R. J., Struck, J., Quinn, P. A., Morgenthaler, N. G., Squire, I. B., Davies, J. E., Bergmann, A., & Ng, L. L. (2007). C-Terminal Provasopressin (Copeptin) as a Novel and Prognostic Marker in Acute Myocardial Infarction. *Leicester Acute Myocardial Infarction Peptide (LAMP) Study*, 115(16), 2103-2110. <https://doi.org/10.1161/circulationaha.106.685503>

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

Knuuti, J., Wijns, W., Saraste, A., Capodanno, D., Barbato, E., Funck-Brentano, C., Prescott, E., Storey, R. F., Deaton, C., Cuisset, T., Agewall, S., Dickstein, K., Edvardsen, T., Escaned, J., Gersh, B. J., Svitil, P., Gilard, M., Hasdai, D., Hatala, R., . . . Group, E. S. C. S. D. (2019). 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes: The Task Force for the diagnosis and management of chronic coronary syndromes of the European Society of Cardiology (ESC). *European Heart Journal*, *41*(3), 407-477.

<https://doi.org/10.1093/eurheartj/ehz425>

Ky, B., French, B., McCloskey, K., Rame, J. E., McIntosh, E., Shahi, P., Dries, D. L., Tang, W. H., Wu, A. H., Fang, J. C., Boxer, R., Sweitzer, N. K., Levy, W. C., Goldberg, L. R., Jessup, M., & Cappola, T. P. (2011). High-sensitivity ST2 for prediction of adverse outcomes in chronic heart failure. *Circ Heart Fail*, *4*(2), 180-187. <https://doi.org/10.1161/circheartfailure.110.958223>

Maisel, A., Mueller, C., Neath, S.-X., Christenson, R. H., Morgenthaler, N. G., McCord, J., Nowak, R. M., Vilke, G., Daniels, L. B., Hollander, J. E., Apple, F. S., Cannon, C., Nagurney, J. T., Schreiber, D., deFilippi, C., Hogan, C., Diercks, D. B., Stein, J. C., Headden, G., . . . Peacock, W. F. (2013). Copeptin Helps in the Early Detection of Patients With Acute Myocardial Infarction: Primary Results of the CHOPIN Trial (Copeptin Helps in the early detection Of Patients with acute myocardial INfarction). *Journal of the American College of Cardiology*, *62*(2), 150-160. <https://doi.org/10.1016/j.jacc.2013.04.011>

Marshall, T., Williams, J., & Williams, K. M. (1991). Electrophoresis of serum isoenzymes and proteins following acute myocardial infarction. *J Chromatogr*, *569*(1-2), 323-345.

[https://doi.org/10.1016/0378-4347\(91\)80236-6](https://doi.org/10.1016/0378-4347(91)80236-6)

McDonagh, T. A., Metra, M., Adamo, M., Gardner, R. S., Baumbach, A., Böhm, M., Burri, H., Butler, J., Čelutkienė, J., Chioncel, O., Cleland, J. G. F., Coats, A. J. S., Crespo-Leiro, M. G., Farmakis, D., Gilard, M., Heymans, S., Hoes, A. W., Jaarsma, T., Jankowska, E. A., . . . Kathrine Skibelund, A. (2021). 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J*, *42*(36), 3599-3726. <https://doi.org/10.1093/eurheartj/ehab368>

McLaurin, M. D., Apple, F. S., Voss, E. M., Herzog, C. A., & Sharkey, S. W. (1997). Cardiac troponin I, cardiac troponin T, and creatine kinase MB in dialysis patients without ischemic heart disease: evidence of cardiac troponin T expression in skeletal muscle. *Clin Chem*, *43*(6 Pt 1), 976-982. <https://doi.org/10.1093/clinchem/43.6.976>

Moussa, I. D., Klein, L. W., Shah, B., Mehran, R., Mack, M. J., Brilakis, E. S., Reilly, J. P., Zoghbi, G., Holper, E., & Stone, G. W. (2013). Consideration of a new definition of clinically relevant myocardial infarction after coronary revascularization: an expert consensus document from the Society for Cardiovascular Angiography and Interventions (SCAI). *J Am Coll Cardiol*, *62*(17), 1563-1570. <https://doi.org/10.1016/j.jacc.2013.08.720>

Neumann, J. T., Twerenbold, R., Ojeda, F., Sorensen, N. A., Chapman, A. R., Shah, A. S. V., Anand, A., Boeddinghaus, J., Nestelberger, T., Badertscher, P., Mokhtari, A., Pickering, J. W., Troughton, R. W., Greenslade, J., Parsonage, W., Mueller-Hennessen, M., Gori, T., Jernberg, T., Morris, N., . . . Blankenberg, S. (2019). Application of High-Sensitivity Troponin in Suspected Myocardial Infarction. *N Engl J Med*, *380*(26), 2529-2540.

<https://doi.org/10.1056/NEJMoa1803377>

Nguyen, T. N., Le, P. X. M., Le, T. X., Nguyen, K. D. A., Nguyen, T. T., Nguyen, T. M., & Tran, V. T. (2020). THE VALUE OF HEART-FATTY ACID BINDING PROTEIN (H-FABP) IN THE EARLY DIAGNOSTIC OF PATIENTS WITH ACUTE MYOCARDIAL INFARCTION. *Journal of the American College of Cardiology*, *75*(11_Supplement_1), 18-18.

[https://doi.org/doi:10.1016/S0735-1097\(20\)30645-8](https://doi.org/doi:10.1016/S0735-1097(20)30645-8)

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

NICE. (2016). Chest pain of recent onset: assessment and diagnosis. <https://www.nice.org.uk/guidance/cg95/chapter/Recommendations>

NICE. (2018, September 12). *Chronic heart failure in adults: diagnosis and management*. <https://www.nice.org.uk/guidance/ng106>

NICE. (2020). High-sensitivity troponin tests for the early rule out of NSTEMI. <https://www.nice.org.uk/guidance/dg40/chapter/1-Recommendations>

O'Connor Robert, E., Al Ali Abdulaziz, S., Brady William, J., Ghaemmaghami Chris, A., Menon, V., Welsford, M., & Shuster, M. (2015). Part 9: Acute Coronary Syndromes. *Circulation*, 132(18_suppl_2), S483-S500. <https://doi.org/10.1161/CIR.0000000000000263>

Pascual-Figal, D. A., & Januzzi, J. L. (2015). The biology of ST2: the International ST2 Consensus Panel. *Am J Cardiol*, 115(7 Suppl), 3b-7b. <https://doi.org/10.1016/j.amjcard.2015.01.034>

Penttila, I., Penttila, K., & Rantanen, T. (2000). Laboratory diagnosis of patients with acute chest pain. *Clin Chem Lab Med*, 38(3), 187-197. <https://doi.org/10.1515/cclm.2000.027>

Ponikowski, P., Voors, A. A., Anker, S. D., Bueno, H., Cleland, J. G. F., Coats, A. J. S., Falk, V., González-Juanatey, J. R., Harjola, V.-P., Jankowska, E. A., Jessup, M., Linde, C., Nihoyannopoulos, P., Parissis, J. T., Pieske, B., Riley, J. P., Rosano, G. M. C., Ruilope, L. M., Ruschitzka, F., . . . Group, E. S. C. S. D. (2016). 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *European Heart Journal*, 37(27), 2129-2200. <https://doi.org/10.1093/eurheartj/ehw128>

Reeder, G., S., Awtry, E., Mahler, S., A. (2022, April 14). *Initial evaluation and management of suspected acute coronary syndrome (myocardial infarction, unstable angina) in the emergency department*. <https://www.uptodate.com/contents/initial-evaluation-and-management-of-suspected-acute-coronary-syndrome-myocardial-infarction-unstable-angina-in-the-emergency-department>

ResponseBio. (2020). POINT OF CARE CARDIAC. <https://responsebio.com/acute-care-diagnostics/cardiovascular/>

Roche. (2020). Roche CARDIAC Trop T Sensitive test (visual). <https://diagnostics.roche.com/global/en/products/params/roche-cardiac-trop-t-sensitive-test-visual.html>

Saggin, L., Gorza, L., Ausoni, S., & Schiaffino, S. (1990). Cardiac troponin T in developing, regenerating and denervated rat skeletal muscle. *Development*, 110(2), 547-554. <https://doi.org/10.1242/dev.110.2.547>

Seino, Y., Ogata, K., Takano, T., Ishii, J., Hishida, H., Morita, H., Takeshita, H., Takagi, Y., Sugiyama, H., Tanaka, T., & Kitaura, Y. (2003). Use of a whole blood rapid panel test for heart-type fatty acid-binding protein in patients with acute chest pain: comparison with rapid troponin T and myoglobin tests. *Am J Med*, 115(3), 185-190. [https://doi.org/10.1016/s0002-9343\(03\)00325-5](https://doi.org/10.1016/s0002-9343(03)00325-5)

Tamis-Holland Jacqueline, E., Jneid, H., Reynolds Harmony, R., Agewall, S., Brilakis Emmanouil, S., Brown Todd, M., Lerman, A., Cushman, M., Kumbhani Dharam, J., Arslanian-Engoren, C., Bolger Ann, F., Beltrame John, F., & null, n. (2019). Contemporary Diagnosis and Management of Patients With Myocardial Infarction in the Absence of Obstructive Coronary

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

Artery Disease: A Scientific Statement From the American Heart Association. *Circulation*, 139(18), e891-e908. <https://doi.org/10.1161/CIR.0000000000000670>

Thygesen, K., Alpert, J. S., Jaffe, A. S., Simoons, M. L., Chaitman, B. R., & White, H. D. (2012). Third Universal Definition of Myocardial Infarction. *Circulation*, 126(16), 2020-2035. <https://doi.org/10.1161/CIR.0b013e31826e1058>

Thygesen, K., Alpert, J. S., & White, H. D. (2007). Universal Definition of Myocardial Infarction. *Circulation*, 116(22), 2634-2653. <https://doi.org/10.1161/circulationaha.107.187397>

Van Nieuwenhoven, F. A., Kleine, A. H., Wodzig, W. H., Hermens, W. T., Kragten, H. A., Maessen, J. G., Punt, C. D., Van Dieijen, M. P., Van der Vusse, G. J., & Glatz, J. F. (1995). Discrimination between myocardial and skeletal muscle injury by assessment of the plasma ratio of myoglobin over fatty acid-binding protein. *Circulation*, 92(10), 2848-2854. <https://doi.org/10.1161/01.CIR.92.10.2848>

Wang, T. J., Wollert, K. C., Larson, M. G., Coglianese, E., McCabe, E. L., Cheng, S., Ho, J. E., Fradley, M. G., Ghorbani, A., Xanthakis, V., Kempf, T., Benjamin, E. J., Levy, D., Vasan, R. S., & Januzzi, J. L. (2012). Prognostic utility of novel biomarkers of cardiovascular stress: the Framingham Heart Study. *Circulation*, 126(13), 1596-1604. <https://doi.org/10.1161/circulationaha.112.129437>

Wang, Y., Tan, X., Gao, H., Yuan, H., Hu, R., Jia, L., Zhu, J., Sun, L., Zhang, H., Huang, L., Zhao, D., Gao, P., & Du, J. (2018). Magnitude of Soluble ST2 as a Novel Biomarker for Acute Aortic Dissection. *Circulation*, 137(3), 259-269. <https://doi.org/10.1161/circulationaha.117.030469>

WHO. (1979). Nomenclature and criteria for diagnosis of ischemic heart disease. Report of the Joint International Society and Federation of Cardiology/World Health Organization task force on standardization of clinical nomenclature. <https://www.ahajournals.org/doi/pdf/10.1161/01.CIR.59.3.607>

Yancy, C. W., Jessup, M., Bozkurt, B., Butler, J., Casey, D. E., Drazner, M. H., Fonarow, G. C., Geraci, S. A., Horwich, T., Januzzi, J. L., Johnson, M. R., Kasper, E. K., Levy, W. C., Masoudi, F. A., McBride, P. E., McMurray, J. J. V., Mitchell, J. E., Peterson, P. N., Riegel, B., . . . Wilkoff, B. L. (2013). 2013 ACCF/AHA Guideline for the Management of Heart Failure: Executive Summary. <https://doi.org/10.1161/CIR.0b013e31829e8807>

Yancy, C. W., Jessup, M., Bozkurt, B., Butler, J., Casey Donald, E., Colvin Monica, M., Drazner Mark, H., Filippatos Gerasimos, S., Fonarow Gregg, C., Givertz Michael, M., Hollenberg Steven, M., Lindenfeld, J., Masoudi Frederick, A., McBride Patrick, E., Peterson Pamela, N., Stevenson Lynne, W., & Westlake, C. (2017). 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation*, 136(6), e137-e161. <https://doi.org/10.1161/CIR.0000000000000509>

Yusuf, S., Collins, R., Lin, L., Sterry, H., Pearson, M., & Sleight, P. (1987). Significance of elevated MB isoenzyme with normal creatine kinase in acute myocardial infarction. *Am J Cardiol*, 59(4), 245-250. [https://doi.org/10.1016/0002-9149\(87\)90793-4](https://doi.org/10.1016/0002-9149(87)90793-4)
Specialty Matched Consultant Advisory Panel review 4/2020

Medical Director review 4/2020

Specialty Matched Consultant Advisory Panel review 4/2021

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “Notification”

Medical Director review 4/2021

Medical Director review 7/2022

Medical Director review 7/2023

Policy Implementation/Update Information

For Policy Titled: Cardiac Biomarkers for Myocardial Infarction

- 1/1/2019 New policy developed. BCBSNC will provide coverage for cardiac biomarkers for myocardial infarction when it is determined to be medically necessary because the criteria and guidelines are met. Medical Director review 1/1/2019. Policy noticed 1/1/2019 for effective date 4/1/2019. (jd)
- 10/01/19 Policy Statement revised to read: BCBSNC will provide coverage for cardiac biomarkers for myocardial infarction when it is determined the medical criteria or reimbursement guidelines noted below are met. Wording revised in the Not Covered section. “Not Medically Necessary” and “investigational” changed to read “Reimbursement is not allowed...” Deleted coding grid. Notification given 10/1/2019 for policy effective date 12/2/2019. (an)
- 4/28/20 Specialty Matched Consultant Advisory Panel review 4/2020. Medical Director review 4/2020. (jd)
- 7/28/20 Reviewed by Avalon 2nd Quarter 2020 CAB. Related policies section added. Policy guidelines and references updated. Medical Director review 7/2020. (jd)
- 5/4/21 Specialty Matched Consultant Advisory Panel review 4/2021. Medical Director review 4/2021. (jd)
- 8/24/21 Reviewed by Avalon 2nd Quarter 2021 CAB. Background, policy guidelines, and references updated. Medical Director review 7/2021 (jd)
- 9/13/22 Reviewed by Avalon 2nd Quarter 2022 CAB. Background, policy guidelines, and references updated. No change to policy statement. Medical Director review 7/2022. (tm)

For Policy Titled: Biomarkers for Myocardial Infarction and Chronic Heart Failure

- 8/15/23 Reviewed by Avalon 2nd Quarter 2023 CAB. Title changed to Biomarkers for Myocardial Infarction and Chronic Heart Failure. Updated Description, Policy Guidelines and References. Policy expanded to include information for the ST2 analysis/assay moved (previously in G2130). When Covered section edited to include repeat troponin testing (moved from previous Note 1), now reads: “For individuals presenting with signs and symptoms of acute coronary syndrome (see Note 1), reimbursement is allowed for **quantitative** measurement of cardiac troponin (troponin T or I) for the diagnosis of myocardial infarction (MI) (when tested at an outpatient facility capable of performing an adequate clinical MI evaluation) up to four times within the first 72 hours following initial presentation.” Not Covered section edited to include new criteria 3 “For all situations, reimbursement is not allowed for qualitative measurement of cardiac troponin (troponin T or I).” and new criteria 5 “For all situations in the outpatient setting, reimbursement is not allowed for analysis of ST2 and/or its isoforms (e.g., Presage ST2).” CPT code 83006 added to Billing/Coding section. Medical Director review 7/2023. **Notification given 8/15/23 for effective date 10/24/23.** (tm)

Biomarkers for Myocardial Infarction and Chronic Heart Failure AHS – G2150 “**Notification**”

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.