

## Corporate Medical Policy

### Psychiatric Partial Hospitalization Programs “NOTIFICATION”

**File Name:** psychiatric\_partial\_hospitalization\_programs  
**Origination:** 03/2024  
**Last Review:** 03/2024

### **Policy Effective 7/1/2024**

#### **Description of Procedure or Service**

---

Psychiatric Partial Hospitalization Programs (PHP) are outpatient care delivery services for Psychiatric Disorders, which must be furnished by or under the supervision of registered or licensed personnel and under the direction of a licensed physician. Partial Hospitalization Programs (PHP) are intended to provide treatment on an outpatient basis, does not include boarding/housing and is intended to provide treatment interventions in a structured setting, with patients returning to their home environments or a community-based setting each day. Partial Hospitalization does not include treatment in a locked unit or restricted access setting.

**Related Policy:**

Psychiatric Intensive Outpatient Programs

***\*\*\*Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.***

#### **Policy**

---

BCBSNC will provide coverage for Psychiatric Partial Hospitalization Programs (PHP) when it is determined to be medically necessary because the medical criteria and guidelines shown below are met.

#### **Benefits Application**

---

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore, member benefit language should be reviewed before applying the terms of this medical policy.

Coverage for services described in this medical policy may be subject to prior authorization by Blue Cross Blue Shield of North Carolina or its designee.

#### **When Psychiatric Partial Hospitalization Programs (PHP) is covered**

---

***Treatment for Psychiatric Partial Hospitalization Programs may be considered medically necessary when members meet ALL the criteria listed below.***

1. If required by state statute, the facility is licensed by the appropriate state agency that approves healthcare facility licensure.
2. There is documentation of drug screens and relevant lab tests upon admission and as clinically indicated.

# Psychiatric Partial Hospitalization Programs “NOTIFICATION”

3. The attending physician is a psychiatrist and responsible for diagnostic evaluation within two days of admission. Thereafter, the physician or physician extender provides an evaluation with documentation as indicated, no less than weekly.
4. After a multidisciplinary assessment, an individualized treatment plan using evidence - based concepts, where applicable, is developed within three days of admission and amended as needed for changes in the individual’s clinical condition. Elements of this plan include, but are not limited to subjects such as identification of key precipitants to current episode of treatment, assessment of psychosocial supports available after discharge, availability of aftercare services in member’s home geographic area, potential need for supportive living placement to continue recovery, consideration of the ability of the member/family/support system to meet financial obligations incurred in the discharge plan, need for services for comorbid medical or substance use conditions, contact with aftercare providers to facilitate an effective transition to lower levels of care, and other issues that affect the likelihood of successful community tenure.
5. Treatment programing includes documentation of at least one individual counseling session weekly or more as clinically indicated.
6. There is documentation that the member is evaluated on each day of the program by a licensed behavioral health practitioner.
7. Licensed behavioral health practitioners supervise all treatment.
8. Mental health and medical services are available 24 hours per day, seven days per week, either on-site or off-site by arrangement.
9. PHP Provides a multidisciplinary treatment program that occurs a minimum 5 days a week and provides a minimum of 20 hours of weekly clinical services to comprehensively address the needs identified in the member’s treatment plan. If the treatment program offers activities that are primarily recreational and diversionary or provide only a level of functional support that does not treat the serious presenting symptoms/problems, BCBSNC does not count these activities in the total hours of treatment delivered.
10. When members are receiving boarding services, during non-program hours the member is allowed the opportunity to:
  - a. Function independently.
  - b. Develop and practice new recovery skills in the real world to prepare for community re-integration and sustained, community-based recovery.
11. There is documentation of a safety plan including access for the member and/or family/support system to professional support outside of program hours.
12. Recent treating providers are contacted by treatment team members to help develop and implement the initial individualized treatment plan within five days of admission.
13. Family participation:
  - a. For adults: Family treatment is being provided at an appropriate frequency. If Family treatment is not rendered, the facility/provider specifically lists the contraindications to Family Therapy.
  - b. For children/adolescents: Family treatment will be provided as part of the treatment plan. If Family treatment is not held, the facility/provider specifically lists the contraindications to Family Therapy. The family/support system assessment will be completed within five days of admission with the expectation that the family is involved in treatment decisions and discharge planning throughout the course of care. Family sessions will occur at least weekly.

# Psychiatric Partial Hospitalization Programs “NOTIFICATION”

- c. Family participation may be conducted via telephonic sessions when there is a significant geographic or other limitation.

## **Admission Criteria:**

### ***Must meet all the following:***

1. A DSM diagnosis is the primary focus of active treatment each program day.
2. There is a reasonable expectation for improvement in the severity of the current condition and behaviors that require a minimum of twenty hours each week to provide treatment, structure and support.
3. The treatment is not primarily social, interpersonal, domiciliary or respite care.
4. The therapeutic supports available in the member’s home community are insufficient to stabilize the member’s current condition and a minimum of twenty hours of treatment each week is required to treat the member’s current condition safely and effectively.
5. The members current condition reflects behavior(s)/psychiatric symptoms that result in functional impairment in 2 areas, including but not limited to:
  - a. potential safety issues for self or others
  - b. primary support
  - c. social/interpersonal
  - d. occupational/educational e. health/medical compliance
6. The member is cognitively capable to actively engage in the recommended treatment plan.
7. This level of care is necessary to provide structure for treatment when at least one of the following exists:
  - a. Clinical documentation supports that the member requires the requested level of care secondary to multiple factors, including but not limited to, medical comorbidity with instability that impairs overall health, concurrent substance use disorder, unstable living situations, a current support system that engages in behaviors that undermine the goals of treatment and adversely affects outcomes, lack of community resources or any other factors that would impact the overall treatment outcome and community tenure.
  - b. After a recent therapeutic trial, the member has a documented history of an inability to adhere to the treatment plan at an intensive\* lower level of care, being non-responsive to treatment or failing to respond to treatment with a reduction in symptom frequency, duration or intensity that triggered the admission. Failure of treatment at a less intensive level of care is not a prerequisite for requiring benefit coverage at a higher level of care.
  - c. The member is at high risk for admission to a higher level of care secondary to multiple recent previous treatments that resulted in unsuccessful stabilization in the community post-discharge.

*\*Note: intensive treatment is defined as at least weekly sessions of individual, family or group counseling*

8. The member needs partial hospitalization because of at least two of the following reasons:
  - a. The member’s condition or stage of recovery requires the need for daily treatment interventions in order to stabilize the clinical condition and acquire the necessary skills to be successful in the next level of care.
  - b. Acute coping skill deficits are significant and require daily assessment and intervention.

# Psychiatric Partial Hospitalization Programs “NOTIFICATION”

- c. A crisis situation is present in social, family, work/school and/or interpersonal relationships which may require daily observation, crisis intervention services, safety planning, problem solving, social services, care coordination, client instruction, support and additional family interventions and other services that may be provided as clinically indicated.

## **Continued Care Criteria:**

***Must meet all the following: (criteria #5 should only be used when the member seeks treatment outside of their home geographic area and #6 only if there are multiple recent admissions)***

1. A DSM diagnosis is the primary focus of active treatment each program day.
2. There is a reasonable expectation for improvement in the severity of the current condition and behaviors that require a minimum of twenty hours each week to provide treatment, structure and support.
3. The treatment is not primarily social, interpersonal, domiciliary or respite care.
4. Family/support system coordination as evidenced by contact with family to discuss current treatment and support needed to transition and maintain treatment at lower levels of care.
5. If a member is receiving treatment outside of their geographic home, discharge planning proactively reflects and mitigates the higher risk of relapse associated with treatment away from home.
6. If a member has a recent history involving multiple treatment attempts with recidivism, the facility develops and implements a treatment plan focused on increasing motivation, readiness for change, practicing new skills to facilitate the development of recovery and other supports to benefit the member in his/her recovery process.
7. The member is displaying increasing motivation, interest in and ability to actively engage in his/her behavioral health treatment, as evidenced by active participation in groups, cooperation with treatment plan, working on assignments, actively developing discharge plan and other markers of treatment engagement. If the member is not displaying increased motivation, there is evidence of active, timely reevaluation and treatment plan modifications to address the current condition.
8. The member’s treatment plan is centered on the alleviation of disabling symptoms and precipitating psychosocial stressors. There is documentation of member progress towards objective, measurable treatment goals that must be met for the member to transition to the next appropriate level of care. If the member is not progressing appropriately or if the member’s condition has worsened, there is evidence of active, timely reevaluation and treatment plan modifications to address the current needs and stabilize the symptoms necessitating the continued stay.
9. The member continues to need partial hospitalization because of at least two of the following:
  - a. The member’s condition or stage of recovery requires the need for daily treatment interventions in order to stabilize the clinical condition and acquire the necessary skills to be successful in the next level of care.
  - b. Acute coping skill deficits are significant and require daily assessment and intervention.
  - c. A crisis situation is present in social, family, work/school and/or interpersonal relationships which may require resources such as crisis intervention services, safety planning, problem solving, social services, care coordination, client instruction, support,

# Psychiatric Partial Hospitalization Programs “NOTIFICATION”

additional family interventions and other services that may be provided as clinically indicated.

10. There is documentation that frequent attempts are made to secure timely access to all current and post-discharge treatment resources and housing (including alternative contingency plans) needed to adequately support timely movement to the next appropriate lower level of care.
11. Despite intensive therapeutic efforts, this level of care is necessary to treat the intensity, frequency and duration of current behaviors and symptoms.

## **When Psychiatric Partial Hospitalization Programs is not covered**

---

Treatment for Psychiatric Partial Hospitalization Programs is considered not medically necessary when members do NOT meet ALL the criteria listed above in the When Treatment is Covered Section.

## **Policy Guidelines**

---

Partial hospital programs (PHPs) provide multidisciplinary behavioral care for 6 to 8 hours per day, 5 to 7 days per week (with patients going home each evening or weekend) and are staffed similarly to the day shift of an inpatient unit.

Symptoms or conditions used to determine the appropriate treatment intensity should be due to the underlying behavioral health diagnosis or represent factors that contribute to destabilization of the underlying diagnosis and are acute in nature or represent a significant worsening over baseline.

Admission to a partial hospital program may be preferable to an intensive outpatient program if daily or near daily management or immediate intervention is necessary. Conditions that may require this intensity of monitoring include medication and comprehensive symptom management; observation and safety planning due to danger to self or others; lack of resiliency and need for repeated reinforcement; extreme mood swings, hopelessness, or isolation with inadequate or unavailable community supports; and substance use monitoring. Immediate intervention may be necessary for crisis situations (e.g., volatile family situations) or when urgent behavioral activation is required (e.g., rapid improvement is necessary to return the individual to vital role functioning)

Impairments in motivation to participate in treatment, limitations to engagement in care, and resistance to change are common in patients with mental health and substance use disorders and may represent a feature of the disease process. A lack of motivation may indicate the need for more intensive services in order to help promote recovery or behavior change. Deficits in motivation or resistance to change should be addressed with therapies designed to enhance motivation to participate in treatment and work toward recovery (e.g., motivational enhancement therapy). The level of motivation and goals/strategies designed to address motivation should be included in the care plan, and a lack of progress toward goals should trigger a reassessment of the care plan (e.g., identification of barriers toward progress, incorporation of new problems that may have developed into the treatment plan, assessment of the appropriateness of initial management strategies), and lead to modification of the plan as appropriate to promote optimal recovery.

## **Billing/Coding/Physician Documentation Information**

---

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at [www.bcbsnc.com](http://www.bcbsnc.com). They are listed in the Category Search on the Medical Policy search page.

*Applicable service codes: H0035*

# Psychiatric Partial Hospitalization Programs “NOTIFICATION”

**Only one (1) unit for PHP on a facility claim is allowed per date of service, as these services are defined as per diem and includes all facility, professional, ancillary, and other services rendered to the member at the site.**

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

## **Scientific Background and Reference Sources**

---

American Association of Community Psychiatrists. Level of Care Utilization System for Psychiatric and Addiction Services (LOCUS). Adult version 20 [Internet] American Association of Community Psychiatrists. 2016 Dec Accessed at: <https://www.communitypsychiatry.org/keystone-programs/locus>

American Medical Association (2020). CPT 2020 Standard Edition. Chicago, IL.

American Psychiatric Association. (2013). Diagnostic and Statistical Manual of Mental Disorders (5th ed., text rev.). Washington, DC.

American Psychiatric Association. Practice guideline for the assessment and treatment of patients with suicidal behaviors 2003; Arlington, VA: American Psychiatric Publishing.

Hunt GE, Siegfried N, Morley K, Brooke-Sumner C, Cleary M. Psychosocial interventions for people with both severe mental illness and substance misuse. Cochrane Database of Systematic Reviews 2019, Issue 12. Art. No.: CD001088. DOI: 10.1002/14651858.CD001088.pub4.

Ijaz S, Davies P, Williams CJ, Kessler D, Lewis G, Wiles N. Psychological therapies for treatment-resistant depression in adults. Cochrane Database of Systematic Reviews 2018, Issue 5. Art. No.: CD010558. DOI: 10.1002/14651858.CD010558.pub2.

Jabbarpour YM, Raney LE. Bridging Transitions of Care from Hospital to Community on the Foundation of Integrated and Collaborative Care. *Focus (Am Psychiatr Publ)*. 2017;15(3):306-315. doi:10.1176/appi.focus.20170017

Kisely S.R., Campbell LA, O'Reilly R. Compulsory community and involuntary outpatient treatment for people with severe mental disorders. Cochrane Database of Systematic Reviews 2017, Issue 3. Art. No.: CD004408. DOI: 10.1002/14651858.CD004408.pub5.

Knable, M. B., Cantrell, C., Meer, A. V., & Levine, E. (2015). The Availability and Effectiveness of Residential Treatment for Persistent Mental Illness. *Psychiatric Annals*, 45(3), 109 -113.

Kemper AR, Maslow GR, Hill S, Namdari B, Allen LaPointe NM, Goode AP, Coeytaux RR, Befus D, Kosinski AS, Bowen SE, McBroom AJ, Lallinger KR, Sanders GD. Attention Deficit Hyperactivity Disorder: Diagnosis and Treatment in Children and Adolescents. Comparative Effectiveness Review No. 203. (Prepared by the Duke University Evidence-based Practice Center under Contract No. 290-2015-00004-I.) AHRQ Publication No. 18-EHC005-EF. Rockville, MD: Agency for Healthcare Research and Quality; January 25, 2018. Posted final reports are located on the Effective Health Care Program search page. DOI: <https://doi.org/10.23970/AHRQEPCCER203>

Leibenluft E. Irritability and Disruptive Mood Dysregulation Disorder. *Journal of the American Academy of Child & Adolescent Psychiatry* 2017, Volume 56, Issue 10, S143 - S144

# Psychiatric Partial Hospitalization Programs “NOTIFICATION”

Machmutow K, Meister R, Jansen A, Kriston L, Watzke B, Härter MC, Liebherz S. Comparative effectiveness of continuation and maintenance treatments for persistent depressive disorder in adults. *Cochrane Database of Systematic Reviews* 2019, Issue 5. Art. No.: CD012855. DOI: 1

Marshall M, Crowther R, Sledge WH, Rathbone J, Soares-Weiser K. Day hospital versus admission for acute psychiatric disorders. *Cochrane Database of Systematic Reviews* 2011, Issue 12. Art. No.: CD004026. DOI: 10.1002/14651858.CD004026.pub2.

McDonagh MS, Dana T, Selph S, Devine EB, Cantor A, Bougatsos C, Blazina I, Grusing S, Fu R, Kopelovich SL, Monroe-DeVita M, Haupt DW. Treatments for Schizophrenia in Adults: A Systematic Review. Comparative Effectiveness Review No. 198. (Prepared by the Pacific Northwest Evidence - based Practice Center under Contract No. 290-2015-00009-I.) AHRQ Publication No. 17(18)-EHC031- EF. Rockville, MD: Agency for Healthcare Research and Quality; October 2017. DOI: <https://doi.org/10.23970/AHRQEPCER198>

McKnight RF, de La Motte de Broöns de Vauvert SJ, Chesney E, Amit BH, Geddes J, Cipriani A. Lithium for acute mania. *Cochrane Database of Systematic Reviews* 2019, Issue 6. Art. No.: CD004048. DOI: 10.1002/14651858.CD004048.pub4.

Michael SS. 2018 AABH Standards and Guidelines, PHP and IOP. Association for Ambulatory Behavioral Healthcare. <https://aabh.org/standards-guidelines/>  
Molyneaux E, Telesia LA, Henshaw C, Boath E, Bradley E, Howard LM. Antidepressants for preventing postnatal depression. *Cochrane Database of Systematic Reviews* 2018, Issue 4. Art. No.: CD004363. DOI: 10.1002/14651858.CD004363.pub3.

Montemagni C, et.al. Second-generation long-acting injectable antipsychotics in schizophrenia: patient functioning and quality of life. *Neuropsychiatric Disease and Treatment* 2016;12 917 – 929

Morant N, et.al. Crisis resolution and home treatment: stakeholders’ views on critical ingredients and implementation in England. *BMC Psychiatry* (2017) 17:254 DOI 10.1186/s12888-017-1421-0

National Institute for Health and Care Excellence (NICE); Bipolar disorder: the assessment and management. London (UK); Published date: 2014 Sep. 58 p. Last updated: April 2018. <https://www.nice.org.uk/guidance/CG185>

North Carolina Department of Health and Human Services; State-Funded Enhanced Mental Health and Substance Abuse Services, October 15, 2023 NCDMA: NC DMA: Title of Policy, Clinical Coverage Policy No. ([ncdhhs.gov](http://ncdhhs.gov))

Olfson M, et.al. Short Term Suicide Risk after Psychiatric Hospital Discharge. *JAMA Psychiatry*. 2016;73(11):1119-1126. doi:10.1001/jamapsychiatry.2016.2035

Oslin D, et.al. VA/DoD Clinical Practice Guideline for Management of Major Depressive Disorder. April 2016. <https://www.healthquality.va.gov/guidelines/MH/mdd/VADoDMDDCPGFINAL82916.pdf>

Ostinelli EG, Brooke-Powney MJ, Li X, Adams CE. Haloperidol for psychosis-induced aggression or agitation (rapid tranquillisation). *Cochrane Database of Systematic Reviews* 2017, Issue 7. Art. No.: CD009377. DOI: 10.1002/14651858.CD009377.pub3.

Ortiz-Orendain J, Castiello-de Obeso S, Colunga-Lozano LE, Hu Y, Maayan N, Adams CE. Antipsychotic combinations for schizophrenia. *Cochrane Database of Systematic Reviews* 2017, Issue 6. Art. No.: CD009005. DOI: 10.1002/14651858.CD009005.pub 2

# Psychiatric Partial Hospitalization Programs “NOTIFICATION”

Ostinelli EG, Hussein M, Ahmed U, Rehman FU, Miramontes K, Adams CE. Risperidone for psychosis-induced aggression or agitation (rapid tranquillisation). *Cochrane Database of Systematic Reviews* 2018, Issue 4. Art. No.: CD009412. DOI: 10.1002/14651858.CD009412.pub2.

Pompoli A, Furukawa TA, Imai H, Tajika A, Efthimiou O, Salanti G. Psychological therapies for panic disorder with or without agoraphobia in adults: a network meta-analysis. *Cochrane Database of Systematic Reviews* 2016, Issue 4. Art. No.: CD011004. DOI: 10.1002/14651858.CD011004.pub2.

Remington G, et.al. Guidelines for the Pharmacotherapy of Schizophrenia in Adults. *The Canadian Journal of Psychiatry / La Revue Canadienne de Psychiatrie* 2017, Vol. 62(9) 604-616 DOI: 10.1177/0706743717720448

Rosser J, Michael S. Partial Hospitalization Programs and Intensive Outpatient Programs. 2021 AABH Standards and Guidelines [Internet] Association for Ambulatory Behavioral Healthcare. 2021 Accessed at: <https://aabh.org/>

Sakinofsky I, et.al. Preventing Suicide among Inpatients. *CanJPsychiatry* 2014;59(3):131–140  
Samara MT, Klupp E, Helfer B, Rothe PH, Schneider-Thoma J, Leucht S. Increasing antipsychotic dose versus switching antipsychotic for non response in schizophrenia. *Cochrane Database of Systematic Reviews* 2018, Issue 5. Art. No.: CD011884. DOI: 10.1002/14651858.CD011884.pub2.

Sampson S, Hosalli P, Furtado VA, Davis JM. Risperidone (depot) for schizophrenia. *Cochrane Database of Systematic Reviews* 2016, Issue 4. Art. No.: CD004161. DOI: 10.1002/14651858.CD004161.pub2.

Sobieraj DM, Baker WL, Martinez BK, Hernandez AV, Coleman CI, Ross JS, Berg KM, Steffens DC. Adverse Effects of Pharmacologic Treatments of Major Depression in Older Adults. *Comparative Effectiveness Review No. 215*. (Prepared by the University of Connecticut Evidence-based Practice Center under Contract No. 290-2015-00012-I.) AHRQ Publication No. 19-EHC011-EF. Rockville, MD: Agency for Healthcare Research and Quality; March 2019. DOI: <https://doi.org/10.23970/AHRQEPCCER215>

Sowers W, et.al. CALOCUS, Child and Adolescent Level of Care Utilization System. American Association of Community Psychiatrists. July 2019. [https://drive.google.com/file/d/0By\\_Xg2nvst9XYjNIWTU0VnAtb11VSnBSQXg2cHU3YUlkMUlZ/view](https://drive.google.com/file/d/0By_Xg2nvst9XYjNIWTU0VnAtb11VSnBSQXg2cHU3YUlkMUlZ/view)

Sowers W, et.al. LOCUS, Level of Care Utilization System. American Association of Community Psychiatrists. December 2016. <https://drive.google.com/file/d/0B89glzXJnn4cV1dESWI2eFEzc3M/view>.

Subramanian S, Völlm BA, Huband N. Clozapine dose for schizophrenia. *Cochrane Database of Systematic Reviews* 2017, Issue 6. Art. No.: CD009555. DOI: 10.1002/14651858.CD009555.pub2

Suomi A, Evans L, Rodgers B, Taplin S, Cowlshaw S. Couple and family therapies for post-traumatic stress disorder (PTSD). *Cochrane Database of Systematic Reviews* 2019, Issue 12. Art. No.: CD011257. DOI: 10.1002/14651858.CD011257.pub2.

The University of South Florida, Florida Medicaid Drug Therapy Management Program sponsored by the Florida Agency for Health Care Administration. 2017-2018 Florida Best Practice Psychotherapeutic Medication Guidelines for Adults (2018).



# Psychiatric Partial Hospitalization Programs “NOTIFICATION”

[http://www.medicaidmentalhealth.org/\\_assets/file/Guidelines/2018Psychotherapeutic%20Medication%20Guidelines%20for%20Adults%20with%20References.pdf](http://www.medicaidmentalhealth.org/_assets/file/Guidelines/2018Psychotherapeutic%20Medication%20Guidelines%20for%20Adults%20with%20References.pdf)

Towbin, Kenneth et al..A Double-Blind Randomized Placebo-Controlled Trial of Citalopram Adjunctive to Stimulant Medication in Youth With Chronic Severe Irritability *Journal of the American Academy of Child & Adolescent Psychiatry* 2019, Volume 59, Issue 3, 350 – 361

Viswanathan M, Kennedy SM, McKeeman J, Christian R, Coker-Schwimmer M, Cook Middleton J, Bann C, Lux L, Randolph C, Forman-Hoffman V. Treatment of Depression in Children and Adolescents: A Systematic Review. Comparative Effectiveness Review No. 224. (Prepared by the RTI International– University of North Carolina at Chapel Hill Evidence-based Practice Center under Contract No. 290-2015-00011-I.) AHRQ Publication No. 20-EHC005-EF. Rockville, MD: Agency for Healthcare Research and Quality; April 2020. DOI: <https://doi.org/10.23970/AHRQEPCER224>

Vita A, Barlati S. The Implementation of Evidence-Based Psychiatric Rehabilitation: Challenges and Opportunities for Mental Health Services. *Front Psychiatry*. 2019; 10:147. Published 2019 Mar 20. doi:10.3389/fpsyt.2019.00147

Weiss A, et.al. Royal Australian and New Zealand College of Psychiatrists Professional Practice Guidelines for the Administration of Electroconvulsive Therapy. *Aust N Z J Psychiatry*. 2019 Jul;609 -623. doi: 10.1177/0004867419839139.

Writing Group and EPPIC National Support Program. Early Psychosis Guidelines. Australian Clinical Guidelines for Early Psychosis, 2nd edition update, 2016, Orygen, The National Centre of Excellence in Youth Mental Health, Melbourne.  
[http://www.ranzcp.org/Files/Resources/Publications/CPG/ClinicalGuidelines-for-Early-Psychosis\\_A-Summary.aspx](http://www.ranzcp.org/Files/Resources/Publications/CPG/ClinicalGuidelines-for-Early-Psychosis_A-Summary.aspx)

Williams T, Hattingh CJ, Kariuki CM, Tromp SA, van Balkom AJ, Ipser JC, Stein DJ. Pharmacotherapy for social anxiety disorder (SAnD). *Cochrane Database of Systematic Reviews* 2017, Issue 10. Art. No.: CD001206. DOI: 10.1002/14651858.CD001206.pub3

Wolraich ML, et.al. Clinical Practice Guideline for the Diagnosis, Evaluation, and Treatment of Attention-Deficit/Hyperactivity Disorder in Children and Adolescents. *PEDIATRICS* Volume 144, number 4, October 2019:e20192528 F

Zaman H, Sampson SJ, Beck ALS, Sharma T, Clay FJ, Spyridi S, Zhao S, Gillies D. Benzodiazepines for psychosis-induced aggression or agitation. *Cochrane Database of Systematic Reviews* 2017, Issue 12. Art. No.: CD003079. DOI: 10.1002/14651858.CD003079.pub4.

Zeanah CH, et.al Practice Parameter for the Assessment and Treatment of Children and Adolescents With Reactive Attachment Disorder and Disinhibited Social Engagement Disorder. *J Am Acad Child Adolesc Psychiatry* 2016;55(11):990–1003.

Zero Suicide in Health and Behavioral Health: <http://zerosuicide.edc.org/>

Medical Director Review 3/2024

---

## Policy Implementation/Update Information

4/1/24 New policy developed. BCBSNC will provide coverage for Psychiatric Partial Hospitalization Programs (PHP) when it is determined to be medically necessary because the medical criteria

# Psychiatric Partial Hospitalization Programs “NOTIFICATION”

and guidelines listed within the policy are met. Medical Director review 3/2024. **Notification given on 4/1/2024 for effective date 7/1/2024.** (tt)

---

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.