

Corporate Medical Policy

Genetic Testing for Hereditary Pancreatitis AHS – M2079

File Name: genetic_testing_for_hereditary_pancreatitis
Origination: 01/2019
Last Review: 07/2024

Description of Procedure or Service

Pancreatitis is defined as inflammation of the pancreas that progresses from acute (AP) (sudden onset; duration <6 months) to recurrent acute (RAP) (>1 episode of acute pancreatitis) to chronic (CP) (duration >6 months) (LaRusch et al., 2014). This recurrent inflammation can lead to total destruction of the pancreas with subsequent pancreatic insufficiency, secondary diabetes, increased risk for pancreatic cancer, and severe unrelenting pain (Ravi Kanth & Nageshwar Reddy, 2014). Hereditary pancreatitis is the early onset form of chronic pancreatitis that is carried in an autosomal dominant pattern with variable penetrance (LaRusch et al., 2012).

Related Policies:

Pancreatic Enzyme Testing for Acute Pancreatitis AHS-G2153
Pancreatic Cancer Risk Testing Using Pancreatic Cyst Fluid AHS-M2114

******Note: This Medical Policy is complex and technical. For questions concerning the technical language and/or specific clinical indications for its use, please consult your physician.***

Policy

BCBSNC will provide coverage for genetic testing for hereditary pancreatitis when it is determined the medical criteria or reimbursement guidelines below are met.

Benefits Application

This medical policy relates only to the services or supplies described herein. Please refer to the Member's Benefit Booklet for availability of benefits. Member's benefits may vary according to benefit design; therefore, member benefit language should be reviewed before applying the terms of this medical policy.

When Genetic Testing for Hereditary Pancreatitis is covered

For individuals under 20 years of age, genetic testing for hereditary pancreatitis (Note 1) is considered medically necessary when at least **one** of the following conditions is met:

- a. For individuals with recurrent (two separate, documented episodes with hyperlipasemia) attacks of acute pancreatitis for which there is no identifiable cause
- b. For individuals with unexplained chronic pancreatitis.
- c. For individuals with a first- or second-degree relative (see Note 2) with a history of recurrent acute pancreatitis, idiopathic chronic pancreatitis, **or** childhood pancreatitis without a known cause.

Genetic Testing for Hereditary Pancreatitis AHS – M2079

- d. For individuals with an unexplained episode of pancreatitis that required hospitalization.

NOTES:

Note 1: For 2 or more gene tests being run on the same platform, please refer to the Laboratory Procedures Medical Policy AHS – R2162.

Note 2: First-degree relatives include parents, full siblings, and children of the individual. Second-degree relatives include grandparents, aunts, uncles, nieces, nephews, grandchildren, and half-siblings of the individual.

When Genetic Testing for Hereditary Pancreatitis is not covered

Genetic testing for hereditary pancreatitis is considered investigational for all other situations not described above.

Policy Guidelines

Background

Pancreatitis is caused by unregulated trypsin activity within the pancreatic acinar cell or pancreatic duct that leads to pancreatic autodigestion and pancreatic inflammation (Lerch & Gorelick, 2000; Whitcomb, 1999). Under acinar cell stress (e.g., hyperstimulation, intracellular hypercalcemia), intracellular trypsinogen is likely converted to trypsin, which activates other digestive enzymes causing injury. Injury releases immune system-activating molecules that cause an initial acute inflammatory response, followed by recruitment of tissue macrophages and activated pancreatic stellate cells. Recurrent injury leads to chronic pancreatitis and fibrosis, mediated by pancreatic stellate cells (Werlin et al., 2015).

Chronic pancreatitis (CP) is a progressive inflammatory disease in which the pancreatic tissue is destroyed over time and replaced by fibrous tissue. The process of fibrosis usually leads to progressive worsening in the structural integrity of the pancreas, changes in arrangement, and composition of the islets, and deformation of the large ducts, eventually resulting in the impairment of both exocrine and endocrine functions (Brock et al., 2013). The annual incidence of the disease has been estimated to be 5-10 per 100,000 persons (Molven et al., 2015). The main symptom of CP is pain; however, it is highly variable in character, frequency, and severity (Mullady et al., 2011; Whitcomb et al., 2008). Therapeutic efforts are mostly aimed at extracting stones and decompressing pancreatic ducts to achieve ideal drainage of the pancreatic duct (Li et al., 2010; Tandan & Nageshwar Reddy, 2013). Genetic risk factors play a larger role in early-onset CP as opposed to late-onset CP. Older adults have a host of environmental and genetic factors that contribute to CP development while hereditary pancreatitis is postulated to make up less than 4% of CP in late-onset groups (Wertheim-Tysarowska et al., 2021).

The etiologies of chronic pancreatitis are classified by the TIGAR-O system into alcoholism, hyperlipidemia, obstructive damage caused by trauma or congenital anomalies, hereditary pancreatitis, autoimmune pancreatitis, and idiopathic (Etemad & Whitcomb, 2001; Sun et al., 2015). The genetic factors listed in TIGAR-O are *PRSSI* (listed as “cationic trypsinogen”), *CFTR*, *SPINK1*, and alpha-1-antitrypsin (listed as “possible”) (Etemad & Whitcomb, 2001). TIGAR-O Version 2 was published in 2019, and lists *PRSSI*, *CFTR*, *SPINK1*, *CTRC*, *CASR*, and *CEL* as genetic factors, as well as some modifier genes such as *CLDN2* (Whitcomb, 2019).

Hereditary pancreatitis (HP) presents as an autosomal dominant chronic pancreatitis with variable penetrance. It mainly develops in childhood (Wertheim-Tysarowska et al., 2021). This variability has been attributed to a genetic predisposition to chronic pancreatitis with the additive effects of environmental and inherited factors. Most genes associated with HP either directly encode components of

Genetic Testing for Hereditary Pancreatitis AHS – M2079

the trypsin system of the exocrine pancreas or are likely to perturb this system indirectly. HP is recognized when pathogenic gene variants of the *PRSSI* gene are found or when acute or chronic pancreatitis develops with a distinct family history (Wertheim-Tysarowska et al., 2021). The phenotype of HP is increased susceptibility to acute pancreatitis, resulting in chronic pancreatitis (including pancreatic fibrosis, chronic pain, maldigestion, and diabetes mellitus) occurring in at least 50%. The risk of pancreatic cancer is also increased (Schwarzenberg, 2023).

Genes Linked to Hereditary Pancreatitis

PRSSI encodes trypsin-1 (cationic trypsinogen), a major pancreatic digestive enzyme. Mutations in *PRSSI* typically result in a trypsin protein that is either prematurely activated or resistant to degradation (LaRusch et al., 2012; Masson et al., 2008), causing autosomal dominant pancreatitis in 60%-100% of families with hereditary pancreatitis (LaRusch & Whitcomb, 2011). “The age of onset for *PRSS-1* related HP ranges from 10 to 12 years” (Hasan et al., 2018).

SPINK1 encodes serine protease inhibitor, Kazal-type 1, a trypsin inhibitor that is upregulated by inflammation (Grendell, 2003). It is not a typical susceptibility gene for acute pancreatitis, but rather a susceptibility gene for the chronic pancreatitis that follows acute pancreatitis.

CTRC encodes chymotrypsin C. Prematurely activated trypsin is destroyed by *CTRC* by acting on the molecule within the calcium-binding loop in the absence of calcium and, therefore, is a crucial candidate gene in the pathogenesis of pancreatitis (Szmola & Sahin-Toth, 2007).

CASR encodes calcium sensing receptor, mutations of which can cause increased calcium ion levels increasing trypsin activation and failed trypsin degradation (Whitcomb, 2004).

CFTR encodes the cystic fibrosis transmembrane conductance protein. Mutations are associated with recurrent acute and chronic pancreatitis since dysfunctional *CFTR* can result in retention of zymogens that can become active and result in pancreatitis (LaRusch & Whitcomb, 2011).

CLDN2 encodes claudin-2, a tight-junction protein that seals the space between epithelial cells. Normally expressed in the proximal pancreatic duct, *CLDN2* is thought to facilitate the transport of water and sodium into the duct to match the chloride and bicarbonate that are actively secreted by pancreatic duct cells through *CFTR*. It is strongly associated with alcohol-related chronic pancreatitis rather than recurrent acute pancreatitis (Ravi Kanth & Nageshwar Reddy, 2014).

CPA1 encodes carboxypeptidase A1; mutated *CPA1* is associated with nonalcoholic chronic pancreatitis, especially with an early age of onset (Witt et al., 2013). Risk for chronic pancreatitis unrelated to trypsin activation appears to be related to endoplasmic reticulum stress from pathogenic *CPA1* variants that alter protein folding, triggering the unfolded protein response.

MYO9B gene and the two tight-junction adaptor genes, ***PARD3*** and ***MAGI2***, have been linked to gastrointestinal permeability. Impairment of the mucosal barrier plays an important role in the pathophysiology of acute pancreatitis (Nijmeijer et al., 2013).

CEL encodes carboxyl-ester lipase, and *CEL* mutations can cause an autosomal dominant syndrome of maturity-onset diabetes of the young (MODY) and exocrine pancreatic dysfunction (Molven et al., 2015).

TRPV6 is a potential novel susceptibility gene for CP that plays a role in epithelial calcium absorption and reabsorption. Variants of this gene also co-occur with pathogenic variants of genes such as *SPINK1* and *CFTR* (Wertheim-Tysarowska et al., 2021).

Genetic Testing for Hereditary Pancreatitis AHS – M2079

Disorder(s)	Genetic Cause(s)	Consequence(s)	Source Citation
Shwachman-Diamond syndrome	<i>SBDS, DNAJC21, EFL1, and SRP54</i>	affect RNA function	(Nelson & Myers, 2008)
Mitochondrial (mt)DNA deletion syndromes, including Kearns-Sayre syndrome (KSS), Pearson syndrome, and progressive external ophthalmoplegia (PEO)	Multiple possible mitochondrial genetic etiologies, including <i>SLC25A4, TWNK, POLG, TYMP, OPA1, RRM2B, DNA2, and MT-TL1,</i>	defective oxidative phosphorylation	(Goldstein & Falk, 2003)
Carboxyl ester lipase (CEL-MODY)	<i>CEL</i>	pancreatic exocrine, endocrine dysfunction, and chronic pancreatitis	(O'Neill et al., 2013)
Johanson-Blizzard syndrome	<i>UBD1</i>	protein synthesis	(Kniffin & McKusick, 2012)

Syndromes that Include Pancreatitis or Pancreatic Insufficiency

Several genes are associated with rare disorders in which pancreatitis or pancreatic insufficiency is part of their phenotype (Durie, 1996; Lerch et al., 2006).

As the number of genes and mutations involved in the onset and progression of pancreatitis becomes higher (Ooi et al., 2010; Walker et al., 2013), the time and cost of screening and sequencing specific genes continues to increase. However, massive parallel sequencing or next generation sequencing (NGS) is becoming standardized (Ballard et al., 2015), and the cost per patient is rapidly dropping (Palermo et

Genetic Testing for Hereditary Pancreatitis AHS – M2079

al., 2016). NGS includes whole genome sequencing, whole exome sequencing (WES) and other methods. Because the cost of WES is now less than the cost of sequencing *CFTR*, use of this technology is becoming an attractive alternative to classic targeted gene sequencing or mutation specific genotyping for a genetic counseling workup (LaRusch et al., 2012). In response to this accelerating development of sequencing techniques, several firms have created genetic panels focusing on hereditary pancreatitis. For example, Invitae offers a six-gene panel (*CASR*, *CFTR*, *CPA1*, *CTRC*, *PRSSI*, *SPINK1*) for chronic pancreatitis (Invitae, 2023). Other firms offering proprietary panels include ARUP Laboratories (4 genes), LabCorp (3 genes), and Ambry (6 genes) (Ambry, 2023; ARUP, 2023; LabCorp, 2023). Still other firms evaluate as many as 12 genes and more (Schwarzenberg, 2023).

Clinical Utility and Validity

Testing for mutations in the *PRSSI*, *SPINK*, and *CFTR* genes is usually done by either direct sequence analysis or next generation sequencing, both of which have high analytic validity. Several studies have evaluated the clinical validity of genetic testing (Applebaum-Shapiro et al., 2001; Ceppa et al., 2013; Poddar et al., 2015; Sultan et al., 2012). One limitation with some studies was lack of inclusion of patients with clinically defined hereditary pancreatitis. Hence, the true clinical sensitivity and specificity of genetic testing in hereditary pancreatitis cannot be accurately determined and needs to be further researched. Similarly, there is a lack of published literature on the clinical utility of testing. Further research is required to evaluate how genetic testing will impact patient management decision and clinical outcomes.

Kumar et al. (2016) sought to characterize and identify risk factors associated with acute recurrent pancreatitis (ARP) and CP in childhood in a multinational cross-sectional study (INSPPIRE). The authors analyzed 301 children with ARP or CP. They found that “At least 1 gene mutation in pancreatitis-related genes was found in 48% of patients with ARP vs 73% of patients with CP. Children with *PRSSI* or *SPINK1* mutations were more likely to present with CP compared with ARP (*PRSSI*: OR = 4.20 and *SPINK1*: OR = 2.30). Obstructive risk factors presented in 33% in both groups, but toxic/metabolic risk factors were more common in children with ARP (21% overall; 26% ARP, 15% CP). They concluded that “The high disease burden in pediatric CP underscores the importance of identifying predisposing factors for progression of ARP to CP in children” (Kumar et al., 2016).

Grabarczyk et al. (2017) also found that *CTRC* variants are strong CP risk factors in pediatric patients. The authors investigated 136 pediatric patients with CP and compared them to 401 controls. They showed that p.Arg254Trp (4.6%) and p.Lys247_Arg254del (5.3%) heterozygous mutations are frequent and significantly associated with CP risk in pediatric patients (odds ratio [OR]=19.1; 95% CI 2.8-160; P=0.001 and OR=5.5; 95% CI 1.6-19.4; P=0.001, respectively). The c.180TT genotype of common p.Gly60Gly variant was found to be a strong and independent CP risk factor (OR=23; 95% CI 7.7-70; P<0.001) with effect size comparable to p.Arg254Trp mutation (Grabarczyk et al., 2017).

Schwarzenberg et al. (2015) evaluated the genetic spectrum of CP. 76 CP patients were examined, and 51 were found to have a genetic risk factor for CP. Of these 51 mutations, 33 were a *PRSSI* mutation, 14 were a *SPINK1* mutation, 11 were a *CFTR* mutation, and 2 were a *CTRC* mutation. The final 25 patients were found to have an obstructive risk factor (Schwarzenberg et al., 2015).

Zou et al. (2018) evaluated the prevalence of four CP-related genes (*SPINK1*, *PRSSI*, *CTRC*, *CFTR*) in Han Chinese patients. The authors performed next-generation sequencing on 1061 patients and 1196 controls. The 1061 patients were further divided into three categories, idiopathic CP (ICP, 715 patients), alcoholic CP (ACP, 206), and smoking-associated CP (SCP, 140). The impact of rare pathogenic variants on age of onset and clinical outcomes was evaluated. Rare pathogenic variants were found in 535 CP patients compared to 71 controls. Mutation positive patients were found to have earlier age of onset as well additional clinical features such as pancreatic stones and diabetes mellitus compared to mutation negative ICP patients. Overall, pathogenic variants were found in 57.1% of ICP patients, compared to 39.8% of ACP patients and 32.1% of SCP patients. The authors concluded that rare pathogenic variants “significantly” influenced age of onset and clinical outcomes of CP (Zou et al., 2018).

Nabi et al. (2020) evaluated 239 children in a prospective study from January 2015 to May 2018 to examine genetic risk factors in children with idiopathic acute recurrent pancreatitis (IARP). Among the enrollees, 85.35% children had IARP, and found that family history of pancreatitis was found among

Genetic Testing for Hereditary Pancreatitis AHS – M2079

4.6% of participants. For specific genes, “mutations/polymorphisms in at least 1 gene were identified in 89.5% (129/144) children including *SPINK1* in 41.9%, *PRSSI* (rs10273639) in 58.2%, *CTRC* in 25.6%, *CTSB* in 54.9%, *CLDN2* in 72.9%, and *CFTR* in 2.3%.” This conveys the overlapping genetic nature of IARP with related genes in HP, making genetic testing important for managing potential disease progression.

Suzuki et al. (2020) investigated the currently understood genetic abnormalities in pancreatitis, and found that “patients with these genetic predispositions [*PRSSI* and *SPINK1* genes], both children and adults, have often been initially diagnosed with idiopathic acute pancreatitis, in approximately 20-50% pediatric cases and 28-0% of adult cases... Patients with chronic pancreatitis (CP) due to *SPINK1* gene mutation and HP patients have a potentially high risk of pancreatic exocrine insufficiency, diabetes mellitus, and of particular importance, pancreatic cancer.” This conveys the continuously emphasized clinical utility of genetic testing to pursue opportunities for counselling and symptom management with disease progression, despite not having gene therapy options for directly targeting HP causing and associated genes (Suzuki et al., 2020).

Weiss et al. (2020) discussed the potential pitfalls from using next generation sequencing (NGS) to diagnose *PRSSI* mutations in chronic pancreatitis. Due to the “high degree of DNA sequence homology (>91%) between *PRSSI* and other members of the trypsinogen multigene family,” there may be erroneous diagnoses of pathologic chronic pancreatitis among patients with benign variants of other *PRSSI*- related genes, like *PRSS2* or *PRSS3P2*. The researchers concluded that sequence homology “can confound the mapping of short NGS reads to a reference genome and lead to technical artefacts.” They recommend “careful clinical evaluation, pretest and post-test genetic counselling and confirmation of NGS test results by Sanger sequencing” to confirm a diagnosis of genetically mutated chronic pancreatitis. This presented the precautions that must be accounted for when utilizing genetic testing for hereditary pancreatitis (Weiss et al., 2020).

Zou et al. (2020) performed whole genome sequencing on a population of 464 Chinese CP patients and on a group of 504 control participants. The *Transient receptor potential cation channel, Subfamily V, Member 6 (TRPV6)* gene was identified as a gene significantly associated with chronic pancreatitis through a “burden test of aggregated rare nonsynonymous variants with a combined annotation dependent depletion score > 20 (p = .020).” In another phase of the study, Sanger sequencing was used to analyze the entire coding sequence and exon/intron boundaries of the *TRPV6* gene. Combining the two phases of the study, the authors identified 25 distinct variants of *TRPV6* and noted that loss-of-function variants were over-represented in the chronic pancreatitis group. The authors concluded that *TRPV6* is likely a novel susceptibility gene for chronic pancreatitis.

Guidelines and Recommendations

Consensus Committees of the European Registry of Hereditary Pancreatic Diseases, the Midwest Multi-Center Pancreatic Study Group and the International Association of Pancreatology

A Consensus Committees of the European Registry of Hereditary Pancreatic Diseases, the Midwest Multi-Center Pancreatic Study Group and the International Association of Pancreatology developed guidelines for genetic testing of the *PRSSI* gene and genetic counseling for HP (Ellis et al., 2001). The recommended indications for symptomatic patients included:

- Recurrent (two separate, documented episodes with hyperlipasemia) attacks of acute pancreatitis for which there is no explanation (anatomical anomalies, ampullary or main pancreatic strictures, trauma, viral infection, gallstones, alcohol, drugs, hyperlipidaemia, etc.)
- Unexplained chronic pancreatitis
- A family history of pancreatitis in a first- or second-degree relative
- Unexplained episode of pancreatitis in a child that required hospitalization

Genetic Testing for Hereditary Pancreatitis AHS – M2079

Predictive (presymptomatic) genetic testing of unaffected relatives is considered more complex. Predictive testing is recommended only for individuals with a first-degree relative with a defined HP gene mutation, and who are over 16 years of age and capable of making an independent and fully informed decision (Ellis et al., 2001).

American Society of Clinical Oncology (ASCO)

The ASCO states that “genetic testing is sometimes considered for patients who develop recurrent pancreatitis at young ages. Genetic testing is available for mutations in the *PRSSI*, *SPINK1*, and *CFTR* genes.” However, the authors note that “it is unknown if mutations in these genes cause an increased risk of pancreatic cancer. The authors believe that other genes may be associated with HP, and studies are ongoing to learn more about this condition” (ASCO, 2021). It is also noted that “Screening for pancreatic cancer is suggested for people known to have HP beginning at age 40, 20 years after the onset of pancreatitis, or 10 years before the youngest pancreatic cancer diagnosis in the family, whichever is earliest. However, the effectiveness of current screening techniques for the early diagnosis of pancreatic cancer is not proven.” Nevertheless, the most commonly used screening tests include magnetic resonance imaging (MRI) and endoscopic ultrasound (EUS) (ASCO, 2021).

American College of Gastroenterology (ACG)

In 2013, the ACG issued guidelines for the management of acute pancreatitis. They include the following recommendation: “genetic testing may be considered in young patients (< 30 years old) if no cause is evident and a family history of pancreatic disease is present (conditional recommendation, low quality of evidence)”. ACG also states that “the role of genetic testing in AP has yet to be determined, but may be useful in patients with more than one family member with pancreatic disease. Individuals with IAP and a family history of pancreatic diseases should be referred for formal genetic counseling” (Tenner et al., 2013).

In 2020, the ACG published clinical guidelines on chronic pancreatitis. There, they state that “In patients with clinical features of CP, a comprehensive review of all risk factors should be performed. This provides information on the underlying mechanisms, identifies both fixed and modifiable risk factors, identifies potential targets for therapies, and provides clinically relevant prognostic information.” As part of that initial approach, they recommend genetic testing in patients “with clinical evidence of a pancreatitis-associated disorder or possible CP [chronic pancreatitis] in which the etiology is unclear, especially in younger patients (strong recommendation, low quality of evidence).” The guideline goes on to state that “at minimum, patients with idiopathic CP should be evaluated for *PRSSI*, *SPINK1*, *CFTR*, and *CTRC* gene mutation analysis...” The guideline mentions that assessment of germline mutations is primarily for prognostic and therapeutic purposes, rather than diagnostic (Gardner et al., 2020).

United European Gastroenterology (UEG)

The United European Gastroenterology published evidence-based guidelines (Lohr et al., 2017) for the diagnosis and therapy of chronic pancreatitis which recommend (Lohr et al., 2017):

“All patients with a family history or early onset disease (<20 years) should be offered genetic testing for associated variants.”

“Genetic screening for every CP patient cannot be recommended since alcohol abuse is the predominant cause of the disease in up to 60% of adult cases.”

“In patients with early onset CP, genetic screening can be offered after informed consent.”

“In patients with alcoholic CP, routine genetic testing cannot be recommended.”

Genetic Testing for Hereditary Pancreatitis AHS – M2079

The working group also noted that “variants in *SPINK1* and *CTRC*, and to a lesser extent, common single-nucleotide polymorphisms (SNPs) in the *PRSSI* and *CLDN2-MORC4* loci, are associated with alcoholic CP” (Lohr et al., 2017).

European Pancreatic Club (EPC) and Hungarian Pancreatic Study Group (HPSG)

The European Pancreatic Club, in collaboration with the Hungarian Pancreatic Study Group organized a consensus guideline meeting on the diagnosis and management of pancreatitis in the pediatric population which state the following:

“Pediatric AP and RAP often develop in the background of genetic susceptibility and genetic testing is warranted in patients with a second episode of idiopathic AP or first episode of idiopathic AP and a family history of AP or CP. Full sequence analysis of *PRSSI*, *SPINK1*, *CTRC*, *CPAI* and *CFTR* gene exons and exon-intron boundaries and testing for the pathogenic CEL hybrid allele are recommended.” The authors go further to mention that “Variants in the *PRSSI* and *CPAI* genes may be associated with a family history of pancreatitis or even autosomal dominant hereditary pancreatitis. Children with a single episode of AP are at risk for developing a second episode. However, genetic testing is cumbersome and expensive. There is usually no therapeutic consequence, but it may assist in long term prognosis” (Parniczky et al., 2018).

“The presence of mutations in the above mentioned genes increases the risk of ARP and CP. Hereditary pancreatitis associated with mutations in *PRSSI*, especially p.R122H, that could considerably increase the risk of pancreatic adenocarcinoma. Knowing the genetic risk factors may not alter the therapy, but it helps to understand the disease's etiological background for the disease and may lead to future targeted investigation” (Parniczky et al., 2018).

Regarding the etiological factors in childhood onset CP, the authors assert that “Genetic variations are the most common risk factors for development of pediatric CP. (*GRADE 1/A, full agreement*) However, other risk factors such as obstruction, autoimmune and toxic and metabolic factors also need to be examined. (*GRADE 2/B, full agreement*)”. Moreover, as “There is an association between CP and cystic fibrosis (CF), therefore a sweat test should be performed to screen for CF as a possible etiological factor in children. (*GRADE 1/A, strong agreement*)” (Parniczky et al., 2018).

International Study Group of Pediatric Pancreatitis: In search for a cuRE (INSPPIRE) Consortium

This group was formed “to collect detailed information on a cohort of children with ARP and CP with the aim to fill gaps in knowledge and improve clinical care”. Their genetic testing-related guidelines are listed below:

- “The search for a genetic cause of ARP or CP should include a sweat chloride test (even if newborn screening for cystic fibrosis (CF) is negative) and *PRSSI* gene mutation testing. Genetic testing for CF should be considered if a sweat test is unable to be performed.”
- “Mutation analysis of the genes *SPINK1*, *CFTR* and *CTRC* may identify risk factors for ARP or CP.”
- “Patients with ARP or CP and a sweat test ≤ 60 mmol/L should have expanded *CFTR* mutation testing done if there is no other identified cause of their pancreatic disease (such as a *PRSSI* mutation or a clear obstructive etiology)” (Garipey et al., 2017).

National Comprehensive Cancer Network (NCCN)

The NCCN notes familial pancreatitis and non-hereditary forms of pancreatitis are both linked with an increased risk of pancreatic cancer. Additionally, chronic pancreatitis is another risk factor for pancreatic cancer. The NCCN specifically lists *PRSSI*, *SPINK1*, and *CFTR* as contributing genes to familial pancreatitis. The approximate increase in risk of pancreatic cancer is somewhere between 26-fold and 87-fold in those with the *PRSSI*, *SPINK1*, and *CFTR* gene mutations (NCCN, 2023).

Genetic Testing for Hereditary Pancreatitis AHS – M2079

International Association of Pancreatology, American Pancreatic Association, Japan Pancreas Society, and European Pancreatic Club

In 2020, the International Association of Pancreatology, the American Pancreatic Association, the Japan Pancreas Society, and European Pancreatic Club released a set of international consensus guidelines on surveillance for pancreatic cancer in the setting of chronic pancreatitis. Though the working group did not explicitly endorse or oppose genetic testing, it was clear that due to the recommendations separated by genetic variants within chronic pancreatitis, genetic testing would become critical for surveillance. With regards to the conditions by which hereditary pancreatitis would warrant surveillance for cancer, the working group stated:

- “The risk of pancreatic cancer in affected individuals with an autosomal dominant history of hereditary pancreatitis due to inherited *PRSSI* mutations is high enough to justify surveillance. *Quality assessment: high; recommendation: strong*”
- “The risk of pancreatic cancer in affected individuals with an autosomal dominant history of hereditary pancreatitis but without *PRSSI* mutations is high enough to justify surveillance. *Quality assessment: moderate; recommendation: weak*”
- “The risk of pancreatic cancer in patients with chronic pancreatitis associated with *SPINK1* p. N34S is not high enough to justify screening or surveillance. *Quality assessment: moderate; recommendation: strong*”
- “The risk of pancreatic cancer in patients with chronic pancreatitis associated with other germline mutations including those of *CFTR*, *CTRC*, *CPAI*, and *CEL*, is not high enough to justify screening or surveillance. *Quality assessment: moderate; recommendation: conditional*” (Greenhalf et al., 2020).

National Institute for Health and Care Excellence (NICE)

NICE updated their guidelines on pancreatitis in December 2020. With regards to genetic testing for hereditary pancreatitis (acute) and patient information, NICE stated the following:

“Give people with pancreatitis, and their family members or carers (as appropriate), written and verbal information on the following, where relevant, as soon as possible after diagnosis:

- pancreatitis and any proposed investigations and procedures, using diagrams
- hereditary pancreatitis, and pancreatitis in children, including specific information on genetic counselling, genetic testing, risk to other family members, and advice on the impact of their pancreatitis on life insurance and travel
- the long-term effects of pancreatitis, including effects on the person's quality of life
- the harm caused to the pancreas by smoking or alcohol.”

For an individual with chronic pancreatitis, NICE recognizes that the cause may not be alcohol-related, but can include “genetic factors; autoimmune disease, in particular IgG4 disease; metabolic causes; [and] structural or anatomical factors” (NICE, 2020).

State and Federal Regulations, as applicable

Food and Drug Administration (FDA)

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

Genetic Testing for Hereditary Pancreatitis AHS – M2079

Billing/Coding/Physician Documentation Information

This policy may apply to the following codes. Inclusion of a code in this section does not guarantee that it will be reimbursed. For further information on reimbursement guidelines, please see Administrative Policies on the Blue Cross Blue Shield of North Carolina web site at www.bcbsnc.com. They are listed in the Category Search on the Medical Policy search page.

Applicable service codes: 81222, 81223, 81224, 81401, 81404, 81405, 81479

BCBSNC may request medical records for determination of medical necessity. When medical records are requested, letters of support and/or explanation are often useful but are not sufficient documentation unless all specific information needed to make a medical necessity determination is included.

Scientific Background and Reference Sources

- Ambry. (2023). Pancreatitis panel. <https://www.ambrygen.com/clinician/genetic-testing/69/exome-and-general-genetics/pancreatitis-panel>
- Applebaum-Shapiro, S. E., Finch, R., Pfutzer, R. H., Hepp, L. A., Gates, L., Amann, S., Martin, S., Ulrich, C. D., 2nd, & Whitcomb, D. C. (2001). Hereditary pancreatitis in North America: the Pittsburgh-Midwest Multi-Center Pancreatic Study Group Study. *Pancreatology*, 1(5), 439-443. <https://doi.org/10.1159/000055844>
- ARUP. (2023). Pancreatitis Panel (CFTR, CTRC, PRSS1, SPINK1), Sequencing. <https://ltd.aruplab.com/Tests/Pub/3004788>
- ASCO. (2021). *Hereditary Pancreatitis*. <https://www.cancer.net/cancer-types/hereditary-pancreatitis>
- Ballard, D. D., Flueckiger, J. R., Fogel, E. L., McHenry, L., Lehman, G. A., Watkins, J. L., Sherman, S., & Cote, G. A. (2015). Evaluating Adults With Idiopathic Pancreatitis for Genetic Predisposition: Higher Prevalence of Abnormal Results With Use of Complete Gene Sequencing. *Pancreas*, 44(1), 116-121. <https://doi.org/10.1097/mpa.0000000000000225>
- Brock, C., Nielsen, L. M., Lelic, D., & Drewes, A. M. (2013). Pathophysiology of chronic pancreatitis. *World J Gastroenterol*, 19(42), 7231-7240. <https://doi.org/10.3748/wjg.v19.i42.7231>
- Ceppa, E. P., Pitt, H. A., Hunter, J. L., Leys, C. M., Zyromski, N. J., Rescorla, F. J., Sandrasegaran, K., Fogel, E. L., McHenry, L. W., Watkins, J. L., Sherman, S., & Lehman, G. A. (2013). Hereditary pancreatitis: endoscopic and surgical management. *J Gastrointest Surg*, 17(5), 847-856; discussion 856-847. <https://doi.org/10.1007/s11605-013-2167-8>
- Durie, P. R. (1996). Inherited and congenital disorders of the exocrine pancreas. *Gastroenterologist*, 4(3), 169-187. <https://www.ncbi.nlm.nih.gov/pubmed/8891682>
- Ellis, I., Lerch, M. M., & Whitcomb, D. C. (2001). Genetic testing for hereditary pancreatitis: guidelines for indications, counselling, consent and privacy issues. *Pancreatology*, 1(5), 405-415. <https://www.karger.com/Article/PDF/55840>
- Etemad, B., & Whitcomb, D. C. (2001). Chronic pancreatitis: diagnosis, classification, and new genetic developments. *Gastroenterology*, 120(3), 682-707. <https://doi.org/10.1053/gast.2001.22586>
- Gardner, T. B., Adler, D. G., Forsmark, C. E., Sauer, B. G., Taylor, J. R., & Whitcomb, D. C. (2020). ACG Clinical Guideline: Chronic Pancreatitis. *American Journal of Gastroenterology*, 115(3). <https://doi.org/10.14309/ajg.0000000000000535>

Genetic Testing for Hereditary Pancreatitis AHS – M2079

- Gariepy, C. E., Heyman, M. B., Lowe, M. E., Pohl, J. F., Werlin, S. L., Wilschanski, M., Barth, B., Fishman, D. S., Freedman, S. D., Giefer, M. J., Gonska, T., Himes, R., Husain, S. Z., Morinville, V. D., Ooi, C. Y., Schwarzenberg, S. J., Troendle, D. M., Yen, E., & Uc, A. (2017). Causal Evaluation of Acute Recurrent and Chronic Pancreatitis in Children: Consensus From the INSPPIRE Group. *J Pediatr Gastroenterol Nutr*, 64(1), 95-103. <https://doi.org/10.1097/mpg.0000000000001446>
- Goldstein, A., & Falk, M. (2003, 01/31/2019). Mitochondrial DNA Deletion Syndromes. University of Washington, WA. <https://www.ncbi.nlm.nih.gov/books/NBK1203/>
- Grabarczyk, A. M., Oracz, G., Wertheim-Tysarowska, K., Kujko, A. A., Wejnarska, K., Kolodziejczyk, E., Bal, J., Koziel, D., Kowalik, A., Gluszek, S., & Rygiel, A. M. (2017). Chymotrypsinogen C Genetic Variants, Including c.180TT, Are Strongly Associated With Chronic Pancreatitis in Pediatric Patients. *J Pediatr Gastroenterol Nutr*, 65(6), 652-657. <https://doi.org/10.1097/mpg.0000000000001767>
- Greenhalf, W., Lévy, P., Gress, T., Rebours, V., Brand, R. E., Pandol, S., Chari, S., Jørgensen, M. T., Mayerle, J., Lerch, M. M., Hegyi, P., Kleeff, J., Castillo, C. F., Isaji, S., Shimosegawa, T., Sheel, A., Halloran, C. M., Garg, P., Takaori, K., . . . Neoptolemos, J. (2020). International consensus guidelines on surveillance for pancreatic cancer in chronic pancreatitis. Recommendations from the working group for the international consensus guidelines for chronic pancreatitis in collaboration with the International Association of Pancreatology, the American Pancreatic Association, the Japan Pancreas Society, and European Pancreatic Club. *Pancreatology*, 20(5), 910-918. <https://doi.org/10.1016/j.pan.2020.05.011>
- Grendell, J. H. (2003). Genetic factors in pancreatitis. *Curr Gastroenterol Rep*, 5(2), 105-109. <https://doi.org/10.1007/s11894-003-0078-7>
- Hasan, A., Moscoso, D. I., & Kastrinos, F. (2018). The Role of Genetics in Pancreatitis. *Gastrointest Endosc Clin N Am*, 28(4), 587-603. <https://doi.org/10.1016/j.giec.2018.06.001>
- Invitae. (2023). Invitae Chronic Pancreatitis Panel. https://www.invitae.com/en/physician/tests/01745/#info-panel-disorders_tested
- Kniffin, C., & McKusick, V. (2012, 09/15/2016). JOHANSON-BLIZZARD SYNDROME; JBS. Johns Hopkins University. <https://omim.org/entry/243800>
- Kumar, S., Ooi, C. Y., Werlin, S., Abu-El-Haija, M., Barth, B., Bellin, M. D., Durie, P. R., Fishman, D. S., Freedman, S. D., Gariepy, C., Giefer, M. J., Gonska, T., Heyman, M. B., Himes, R., Husain, S. Z., Lin, T. K., Lowe, M. E., Morinville, V., Palermo, J. J., . . . Uc, A. (2016). Risk Factors Associated With Pediatric Acute Recurrent and Chronic Pancreatitis: Lessons From INSPPIRE. *JAMA Pediatr*, 170(6), 562-569. <https://doi.org/10.1001/jamapediatrics.2015.4955>
- LabCorp. (2023). Pancreatitis: Three-gene Profile (PRSS1, SPINK1, CFTR) (Full Gene Sequencing). <https://www.labcorp.com/tests/252794/pancreatitis-three-gene-profile-i-prss1-spink1-cfr-i-full-gene-sequencing>
- LaRusch, J., Barmada, M. M., Solomon, S., & Whitcomb, D. C. (2012). Whole exome sequencing identifies multiple, complex etiologies in an idiopathic hereditary pancreatitis kindred. *Jop*, 13(3), 258-262. <https://pubmed.ncbi.nlm.nih.gov/22572128/>
- LaRusch, J., Solomon, S., & Whitcomb, D. C. (2014). Pancreatitis Overview [Text]. *GENEReviews*. <https://www.ncbi.nlm.nih.gov/books/NBK190101/>
- LaRusch, J., & Whitcomb, D. C. (2011). Genetics of pancreatitis. *Curr Opin Gastroenterol*, 27(5), 467-474. <https://doi.org/10.1097/MOG.0b013e328349e2f8>

Genetic Testing for Hereditary Pancreatitis AHS – M2079

- Lerch, M. M., & Gorelick, F. S. (2000). Early trypsinogen activation in acute pancreatitis. *Med Clin North Am*, 84(3), 549-563, viii. [https://doi.org/10.1016/S0025-7125\(05\)70239-X](https://doi.org/10.1016/S0025-7125(05)70239-X)
- Lerch, M. M., Zenker, M., Turi, S., & Mayerle, J. (2006). Developmental and metabolic disorders of the pancreas. *Endocrinol Metab Clin North Am*, 35(2), 219-241, vii. <https://doi.org/10.1016/j.ecl.2006.02.004>
- Li, Z. S., Wang, W., Liao, Z., Zou, D. W., Jin, Z. D., Chen, J., Wu, R. P., Liu, F., Wang, L. W., Shi, X. G., Yang, Z., & Wang, L. (2010). A long-term follow-up study on endoscopic management of children and adolescents with chronic pancreatitis. *Am J Gastroenterol*, 105(8), 1884-1892. <https://doi.org/10.1038/ajg.2010.85>
- Lohr, J. M., Dominguez-Munoz, E., Rosendahl, J., Besselink, M., Mayerle, J., Lerch, M. M., Haas, S., Akisik, F., Kartalis, N., Iglesias-Garcia, J., Keller, J., Boermeester, M., Werner, J., Dumonceau, J. M., Fockens, P., Drewes, A., Ceyhan, G., Lindkvist, B., Drenth, J., . . . Bruno, M. (2017). United European Gastroenterology evidence-based guidelines for the diagnosis and therapy of chronic pancreatitis (HaPanEU). *United European Gastroenterol J*, 5(2), 153-199. <https://doi.org/10.1177/2050640616684695>
- Masson, E., Le Marechal, C., Delcenserie, R., Chen, J. M., & Ferec, C. (2008). Hereditary pancreatitis caused by a double gain-of-function trypsinogen mutation. *Hum Genet*, 123(5), 521-529. <https://doi.org/10.1007/s00439-008-0508-6>
- Molven, A., Njolstad, P. R., & Weiss, F. U. (2015). Lipase gene fusion: a new route to chronic pancreatitis. In *Oncotarget* (Vol. 6, pp. 30443-30444). <https://doi.org/10.18632/oncotarget.5454>
- Mullady, D. K., Yadav, D., Amann, S. T., O'Connell, M. R., Barmada, M. M., Elta, G. H., Scheiman, J. M., Wamsteker, E. J., Chey, W. D., Korneffel, M. L., Weinman, B. M., Slivka, A., Sherman, S., Hawes, R. H., Brand, R. E., Burton, F. R., Lewis, M. D., Gardner, T. B., Gelrud, A., . . . Anderson, M. A. (2011). Type of pain, pain-associated complications, quality of life, disability and resource utilisation in chronic pancreatitis: a prospective cohort study. *Gut*, 60(1), 77-84. <https://doi.org/10.1136/gut.2010.213835>
- Nabi, Z., Talukdar, R., Venkata, R., Aslam, M., Shava, U., & Reddy, D. N. (2020). Genetic Evaluation of Children with Idiopathic Recurrent Acute Pancreatitis. *Dig Dis Sci*, 65(10), 3000-3005. <https://doi.org/10.1007/s10620-019-06026-2>
- NCCN. (2023, December 13, 2023). *NCCN Clinical Practice Guidelines in Oncology - Pancreatic Adenocarcinoma*
- Nelson, A., & Myers, K. (2008, 10/18/2018). Shwachman-Diamond Syndrome. University of Washington, Seattle. <https://www.ncbi.nlm.nih.gov/books/NBK1756/>
- NICE. (2020, December 16). Pancreatitis. Retrieved April 22 from <https://www.nice.org.uk/guidance/ng104/chapter/Recommendations>
- Nijmeijer, R. M., van Santvoort, H. C., Zhernakova, A., Teller, S., Scheiber, J. A., de Kovel, C. G., Besselink, M. G., Visser, J. T., Lutgendorff, F., Bollen, T. L., Boermeester, M. A., Rijkers, G. T., Weiss, F. U., Mayerle, J., Lerch, M. M., Gooszen, H. G., Akkermans, L. M., & Wijnenga, C. (2013). Association analysis of genetic variants in the myosin IXB gene in acute pancreatitis. *PLoS One*, 8(12), e85870. <https://doi.org/10.1371/journal.pone.0085870>
- O'Neill, M., Stumpf, A., & McKusick, V. (2013, 09/10/2013). MATURITY-ONSET DIABETES OF THE YOUNG, TYPE 8, WITH EXOCRINE DYSFUNCTION; MODY8. Johns Hopkins University. <https://omim.org/entry/609812>

Genetic Testing for Hereditary Pancreatitis AHS – M2079

- Ooi, C. Y., Gonska, T., Durie, P. R., & Freedman, S. D. (2010). Genetic testing in pancreatitis. *Gastroenterology*, 138(7), 2202-2206, 2206.e2201. <https://doi.org/10.1053/j.gastro.2010.04.022>
- Palermo, J. J., Lin, T. K., Hornung, L., Valencia, C. A., Mathur, A., Jackson, K., Fei, L., & Abu-El-Haija, M. (2016). Genophenotypic Analysis of Pediatric Patients With Acute Recurrent and Chronic Pancreatitis. *Pancreas*, 45(9), 1347-1352. <https://doi.org/10.1097/mpa.0000000000000655>
- Parniczky, A., Abu-El-Haija, M., Husain, S., Lowe, M., Oracz, G., Sahin-Toth, M., Szabo, F. K., Uc, A., Wilschanski, M., Witt, H., Czako, L., Grammatikopoulos, T., Rasmussen, I. C., Sutton, R., & Hegyi, P. (2018). EPC/HPSG evidence-based guidelines for the management of pediatric pancreatitis. *Pancreatology*, 18(2), 146-160. <https://doi.org/10.1016/j.pan.2018.01.001>
- Poddar, U., Yachha, S. K., Mathias, A., & Choudhuri, G. (2015). Genetic predisposition and its impact on natural history of idiopathic acute and acute recurrent pancreatitis in children. *Dig Liver Dis*, 47(8), 709-714. <https://doi.org/10.1016/j.dld.2015.04.012>
- Ravi Kanth, V., & Nageshwar Reddy, D. (2014). Genetics of acute and chronic pancreatitis: An update. *World J Gastrointest Pathophysiol*, 5(4), 427-437. <https://doi.org/10.4291/wjgp.v5.i4.427>
- Schwarzenberg, S. J. (2023, September 13, 2023). *Pancreatitis associated with genetic risk factors*. UpToDate, Inc. <https://www.uptodate.com/contents/pancreatitis-associated-with-genetic-risk-factors>
- Schwarzenberg, S. J., Bellin, M., Husain, S. Z., Ahuja, M., Barth, B., Davis, H., Durie, P. R., Fishman, D. S., Freedman, S. D., Garipey, C. E., Giefer, M. J., Gonska, T., Heyman, M. B., Himes, R., Kumar, S., Morinville, V. D., Lowe, M. E., Nuehring, N. E., Ooi, C. Y., . . . Uc, A. (2015). Pediatric chronic pancreatitis is associated with genetic risk factors and substantial disease burden. *J Pediatr*, 166(4), 890-896.e891. <https://doi.org/10.1016/j.jpeds.2014.11.019>
- Sultan, M., Werlin, S., & Venkatasubramani, N. (2012). Genetic prevalence and characteristics in children with recurrent pancreatitis. *J Pediatr Gastroenterol Nutr*, 54(5), 645-650. <https://doi.org/10.1097/mpg.0b013e31823f0269>
- Sun, C., Liu, M. Y., Liu, X. G., Hu, L. H., Xia, T., Liao, Z., & Li, Z. S. (2015). Serine Protease Inhibitor Kazal Type 1 (SPINK1) c.194+2T > C Mutation May Predict Long-term Outcome of Endoscopic Treatments in Idiopathic Chronic Pancreatitis. *Medicine (Baltimore)*, 94(47), e2046. <https://doi.org/10.1097/md.0000000000002046>
- Suzuki, M., Minowa, K., Nakano, S., Isayama, H., & Shimizu, T. (2020). Genetic Abnormalities in Pancreatitis: An Update on Diagnosis, Clinical Features, and Treatment. *Diagnostics (Basel, Switzerland)*, 11(1), 31. <https://doi.org/10.3390/diagnostics11010031>
- Szmola, R., & Sahin-Toth, M. (2007). Chymotrypsin C (caldecrin) promotes degradation of human cationic trypsin: identity with Rinderknecht's enzyme Y. *Proc Natl Acad Sci U S A*, 104(27), 11227-11232. <https://doi.org/10.1073/pnas.0703714104>
- Tandan, M., & Nageshwar Reddy, D. (2013). Endotherapy in chronic pancreatitis. *World J Gastroenterol*, 19(37), 6156-6164. <https://doi.org/10.3748/wjg.v19.i37.6156>
- Tenner, S., Baillie, J., DeWitt, J., & Vege, S. S. (2013). American College of Gastroenterology guideline: management of acute pancreatitis. *Am J Gastroenterol*, 108(9), 1400-1415; 1416. <https://doi.org/10.1038/ajg.2013.218>
- Walker, N. F., Warren, O. J., Gawn, L., & Jiao, L. R. (2013). The role of genetic testing in management of hereditary chronic pancreatitis. *JRSM Short Rep*, 4(1), 6. <https://doi.org/10.1258/shorts.2012.012071>

Genetic Testing for Hereditary Pancreatitis AHS – M2079

- Weiss, F. U., Laemmerhirt, F., & Lerch, M. M. (2020). Next generation sequencing pitfalls in diagnosing trypsinogen (PRSS1) mutations in chronic pancreatitis. *Gut*. <https://doi.org/10.1136/gutjnl-2020-322864>
- Werlin, S., Konikoff, F. M., Halpern, Z., Barkay, O., Yerushalmi, B., Broide, E., Santo, E., Shamir, R., Shaoul, R., Shteyer, E., Yaakov, Y., Cohen, M., Kerem, E., Ruzniewski, P., Masson, E., Ferec, C., & Wilschanski, M. (2015). Genetic and electrophysiological characteristics of recurrent acute pancreatitis. *J Pediatr Gastroenterol Nutr*, 60(5), 675-679. <https://doi.org/10.1097/mpg.0000000000000623>
- Wertheim-Tysarowska, K., Oracz, G., & Rygiel, A. M. (2021). Genetic Risk Factors in Early-Onset Nonalcoholic Chronic Pancreatitis: An Update. *Genes*, 12(5), 785. <https://doi.org/10.3390/genes12050785>
- Whitcomb, D. C. (1999). Early trypsinogen activation in acute pancreatitis. *Gastroenterology*, 116(3), 770-772. [https://www.gastrojournal.org/article/S0016-5085\(99\)70205-2/fulltext](https://www.gastrojournal.org/article/S0016-5085(99)70205-2/fulltext)
- Whitcomb, D. C. (2004). Mechanisms of disease: Advances in understanding the mechanisms leading to chronic pancreatitis. *Nat Clin Pract Gastroenterol Hepatol*, 1(1), 46-52. <https://doi.org/10.1038/ncpgasthep0025>
- Whitcomb, D. C. (2019). Pancreatitis: TIGAR-O Version 2 Risk/Etiology Checklist With Topic Reviews, Updates, and Use Primers. *Clin Transl Gastroenterol*, 10(6), e00027. <https://doi.org/10.14309/ctg.0000000000000027>
- Whitcomb, D. C., Yadav, D., Adam, S., Hawes, R. H., Brand, R. E., Anderson, M. A., Money, M. E., Banks, P. A., Bishop, M. D., Baillie, J., Sherman, S., DiSario, J., Burton, F. R., Gardner, T. B., Amann, S. T., Gelrud, A., Lo, S. K., DeMeo, M. T., Steinberg, W. M., . . . Barmada, M. M. (2008). Multicenter approach to recurrent acute and chronic pancreatitis in the United States: the North American Pancreatitis Study 2 (NAPS2). *Pancreatology*, 8(4-5), 520-531. <https://doi.org/10.1159/000152001>
- Witt, H., Beer, S., Rosendahl, J., Chen, J. M., Chandak, G. R., Masamune, A., Bence, M., Szmola, R., Oracz, G., Macek, M., Jr., Bhatia, E., Steigenberger, S., Lasher, D., Buhler, F., Delaporte, C., Tebbing, J., Ludwig, M., Pilsak, C., Saum, K., . . . Sahin-Toth, M. (2013). Variants in CPA1 are strongly associated with early onset chronic pancreatitis. *Nat Genet*, 45(10), 1216-1220. <https://doi.org/10.1038/ng.2730>
- Zou, W. B., Tang, X. Y., Zhou, D. Z., Qian, Y. Y., Hu, L. H., Yu, F. F., Yu, D., Wu, H., Deng, S. J., Lin, J. H., Zhao, A. J., Zhao, Z. H., Wu, H. Y., Zhu, J. H., Qian, W., Wang, L., Xin, L., Wang, M. J., Wang, L. J., . . . Liao, Z. (2018). SPINK1, PRSS1, CTSC, and CFTR Genotypes Influence Disease Onset and Clinical Outcomes in Chronic Pancreatitis. *Clin Transl Gastroenterol*, 9(11), 204. <https://doi.org/10.1038/s41424-018-0069-5>
- Zou, W. B., Wang, Y. C., Ren, X. L., Wang, L., Deng, S. J., Mao, X. T., Li, Z. S., & Liao, Z. (2020). TRPV6 variants confer susceptibility to chronic pancreatitis in the Chinese population. *Hum Mutat*, 41(8), 1351-1357. <https://doi.org/10.1002/humu.24032>
- Specialty Matched Consultant Advisory Panel review 7/2019
- Medical Director review 7/2019
- Specialty Matched Consultant Advisory Panel review 7/2020

Genetic Testing for Hereditary Pancreatitis AHS – M2079

Medical Director review 7/2020

Specialty Matched Consultant Advisory Panel review 7/2021

Medical Director review 7/2021

Medical Director review 7/2022

Medical Director review 7/2023

Medical Director review 7/2024

Policy Implementation/Update Information

- 1/1/2019 BCBSNC will provide coverage for genetic testing for hereditary pancreatitis when it is determined to be medically necessary because the medical criteria and guidelines are met. Medical Director review 1/1/2019. Policy noticed 1/1/2019 for effective date 4/1/2019 (jd)
- 8/13/2019 Specialty Matched Consultant Advisory Panel review 7/2019. Medical Director review 7/2019. (jd)
- 9/10/2019 Reviewed by Avalon 2nd Quarter 2019 CAB. Related Policies added to Description section. Minor revision to When Covered section; removed “Unexplained pancreatitis with” from item 1c. Policy guidelines and references updated. Code table removed from Billing/Coding section. Medical Director review 8/2019. (jd)
- 10/29/19 Wording in the Policy, When Covered, and/or Not Covered section(s) changed from Medical Necessity to Reimbursement language, where needed. (hb)
- 7/28/20 Reviewed by Avalon 2nd Quarter 2020 CAB. Note 1 added to When Covered section. Policy guidelines and references updated. Specialty Matched Consultant Advisory Panel review 7/2020. Medical Director review 7/2020. (jd)
- 8/24/21 Reviewed by Avalon 2nd Quarter 2021 CAB. Policy guidelines and references updated. Specialty Matched Consultant Advisory Panel review 7/2021. Medical Director review 7/2021. (jd)
- 9/13/22 Reviewed by Avalon 2nd Quarter 2022 CAB. Description, policy guidelines, background and references updated. When Covered section edited for clarity. No change to policy statement. Medical Director review 7/2022. (tm)
- 8/15/23 Reviewed by Avalon 2nd Quarter 2023 CAB. Updated the background, guidelines and recommendations, and evidence-based scientific references. When covered section edited for clarity, no changes to policy statement. Medical Director review 7/2023. (rp)
- 9/4/24 Reviewed by Avalon 2nd Quarter 2024 CAB. Updated Description, Policy Guidelines, and References. Note 1 under When Covered section updated to reflect changes to definition of a genetic panel within R2162. Now reads: “Note 1: For 2 or more gene tests being run on the same platform, please refer to Laboratory Procedures Medical Policy AHS - R2162.” Medical Director review 7/2024. (tm)

Medical policy is not an authorization, certification, explanation of benefits or a contract. Benefits and eligibility are determined before medical guidelines and payment guidelines are applied. Benefits are determined by the group contract and subscriber certificate that is in effect at the time services are rendered. This document is solely provided for informational

Genetic Testing for Hereditary Pancreatitis AHS – M2079

purposes only and is based on research of current medical literature and review of common medical practices in the treatment and diagnosis of disease. Medical practices and knowledge are constantly changing and BCBSNC reserves the right to review and revise its medical policies periodically.